

## **ПРЕПРИНТЫ ПОМИ РАН**

### **ГЛАВНЫЙ РЕДАКТОР**

**С.В. Кисляков**

### **РЕДКОЛЛЕГИЯ**

**В.М.Бабич, Н.А.Вавилов, А.М.Вершик, М.А.Всемирнов, А.И.Генералов, И.А.Ибрагимов,  
Л.Ю.Колотилина, Б.Б.Лурье, Ю.В.Матиясевич, Н.Ю.Нецветаев, С.И.Репин, Г.А.Серегин**

**Учредитель: Федеральное государственное бюджетное учреждение науки  
Санкт-Петербургское отделение Математического института  
им. В. А. Стеклова Российской академии наук**

**Свидетельство о регистрации средства массовой информации: ЭЛ №ФС 77-33560 от 16  
октября 2008 г. Выдано Федеральной службой по надзору в сфере связи и массовых  
коммуникаций**

**Контактные данные: 191023, г. Санкт-Петербург, наб. реки Фонтанки, дом 27**

**телефоны: (812)312-40-58; (812) 571-57-54**

**e-mail: [admin@pdmi.ras.ru](mailto:admin@pdmi.ras.ru)**

**<http://www.pdmi.ras.ru/preprint/>**

**Заведующая информационно-издательским сектором Симонова В.Н**

$L_p$ -THEORY OF FREE BOUNDARY PROBLEMS  
OF MAGNETOHYDRODYNAMICS IN MULTI-CONNECTED DOMAINS

V. A. Solonnikov

*Dedicated to the memory of O. A. Ladyzhenskaya*

St.Petersburg Department of V. A. Steklov Institute  
of Mathematics of Russian Academy of Sciences

[solonnik@pdmi.ras.ru](mailto:solonnik@pdmi.ras.ru)

September 3, 2013

ABSTRACT:

We develop the  $L_p$ -theory of solvability of free boundary problems of magnetohydrodynamics of viscous incompressible fluids in multi-connected domains constructed in the paper [1] for  $p = 2$ . The case of simply connected domains is studied in [2,3].

**Key words:** free boundaries, Sobolev spaces, magnetohydrodynamics, divergence free vector fields

The paper was supported by the RFBR grant 11-01-00-324.

## ГЛАВНЫЙ РЕДАКТОР

С. В. Кисляков

## РЕДКОЛЛЕГИЯ

В. М. Бабич, А. Н. Бородин, Н. А. Вавилов, А. М. Вершик, М. А. Всемиров, А. И. Генералов, А. Ю. Зайцев, И. А. Ибрагимов, Л. Ю. Колотилина, Г. В. Кузьмина, П. П. Кулиш, М. А. Лифшиц, Б. Б. Лурье, Ю. В. Матиясевич, Н. Ю. Нечетаев, С. И. Репин, Г. А. Серегин, В. Н. Судаков, О. М. Фоменко

# 1 Introduction.

As in [1], we consider the problem of finding a bounded variable domain  $\Omega_{1t} \subset \mathbb{R}^3$  with the boundary  $\Gamma_t$ ,  $t > 0$ , filled with the fluid, together with the vector fields of velocity  $\mathbf{v}(x, t)$ , magnetic and electric fields  $\mathbf{H}(x, t)$ ,  $\mathbf{E}(x, t)$  and the pressure function  $p(x, t)$ . We assume that the fluid is subject to the mass forces, capillary force at the free boundary  $\Gamma_t$  and forces due to the presence of the magnetic and electric fields that are generated by the electric current  $\mathbf{j}(x, t)$  given in a fixed domain  $\Omega_3$  that is bounded away from  $\Omega_{1t}$ . The function  $\mathbf{j}(x, t)$  satisfies the condition

$$\mathbf{j}(x, t) \cdot \mathbf{n}(x) = 0, \quad x \in \partial\Omega_3 \equiv S_3 \quad (1.1)$$

and vanishes outside  $\Omega_3$ . Both  $\Omega_{1t}$  and  $\Omega_3$  are surrounded by a vacuum region  $\Omega_{2t}$ . The domain  $\Omega = \bar{\Omega}_{1t} \cup \bar{\Omega}_3 \cup \Omega_{2t}$  is bounded by a perfectly conducting surface  $S$ . The governing equations are the Navier-Stokes equations with the magnetic field  $\mathbf{H}$  and the Maxwell equations without displacement current (i.e., without the derivative  $\mathbf{E}_t$ ) - see [4.5]:

$$\begin{cases} \mathbf{v}_t + (\mathbf{v} \cdot \nabla)\mathbf{v} - \nabla \cdot T(\mathbf{v}, p) - \nabla \cdot T_M(\mathbf{H}) = \mathbf{f}(x, t), \\ \nabla \cdot \mathbf{v}(x, t) = 0, \quad x \in \Omega_{1t}, \quad t > 0, \end{cases} \quad (1.2)$$

$$\begin{cases} \mu \mathbf{H}_t = -\text{rot} \mathbf{E}, \quad \nabla \cdot \mathbf{H} = 0, \quad x \in \Omega_{1t} \cup \Omega_{2t} \cup \Omega_3, \\ \text{rot} \mathbf{H} = \alpha_1(\mathbf{E} + \mu_1(\mathbf{v} \times \mathbf{H})), \quad x \in \Omega_{1t}, \quad t > 0, \\ \text{rot} \mathbf{H} = \alpha_3 \mathbf{E} + \mathbf{j}(x, t), \quad x \in \Omega_3, \\ \text{rot} \mathbf{H} = 0, \quad \nabla \cdot \mathbf{H} = 0, \quad \nabla \cdot \mathbf{E} = 0, \quad x \in \Omega_{2t}. \end{cases} \quad (1.3)$$

These equations are completed by the following boundary and initial conditions:

$$\begin{cases} (T(\mathbf{v}, p) + [T_M(\mathbf{H})])\mathbf{n} = \sigma \mathbf{n} H, \quad V_n = \mathbf{v} \cdot \mathbf{n}, \quad x \in \Gamma_t, \\ \mathbf{n}_t[\mu \mathbf{H}] + [\mathbf{n}_x \times \mathbf{E}] = 0, \quad x \in \Gamma_t, \\ [\mu \mathbf{H} \cdot \mathbf{n}] = 0, \quad [\mathbf{H}_\tau] = 0, \quad x \in \Gamma_t \cup S_3, \quad [\mathbf{E}_\tau] = 0, \quad x \in S_3, \\ \mathbf{H} \cdot \mathbf{n} = 0, \quad \mathbf{E}_\tau = 0, \quad x \in S, \\ \mathbf{v}(x, 0) = \mathbf{v}_0(x), \quad x \in \Omega_{10}, \quad \mathbf{H}(x, 0) = \mathbf{H}_0(x), \quad x \in \Omega_{10} \cup \Omega_{20} \cup \Omega_3. \end{cases} \quad (1.4)$$

Here  $T(\mathbf{v}, p)$  is the viscous stress tensor:  $T(\mathbf{v}, p) = -pI + \nu S(\mathbf{v})$ ,  $S(\mathbf{v}) = \nabla \mathbf{v} + (\nabla \mathbf{v})^T$  is the doubled rate-of-strain tensor,  $T_M(\mathbf{H}) = \mu(\mathbf{H} \otimes \mathbf{H} - \frac{1}{2}|\mathbf{H}|^2 I)$ , is the magnetic stress tensor,  $\mu$  and  $\alpha$  are piece-wise constant functions equal to  $\mu_i$  and  $\alpha_i$  in  $\Omega_{it}$ ,  $\mu = \mu_3$  in  $\Omega_3$ ,  $\alpha = 0$  in  $\Omega_{2t}$ ,  $\mathbf{n}$  is the normal to  $\Gamma_t$ ,  $S_3$ ,  $S$ , exterior with respect to  $\Omega_{1t}$ ,  $\Omega_3$ ,  $\Omega$ ,  $V_n$  is the velocity of evolution of  $\Gamma_t$  in the direction  $\mathbf{n}$ ,  $\mathbf{H}_\tau = \mathbf{H} - \mathbf{n}(\mathbf{n} \cdot \mathbf{H})$  is the tangential component of  $\mathbf{H}$ ,  $H$  is the doubled mean curvature of  $\Gamma_t$  negative for convex domains. The parameters  $\nu$ ,  $\mu_i$ ,  $\alpha_i$ ,  $\sigma$  are positive constants and  $[u]$  is the jump of the function  $u(x)$ ,  $x \in \Omega_i$ ,  $i = 1, 2, 3$ , on  $\Gamma_t$  or  $S_3$ . We set  $[u]|_{\Gamma_t} = u^{(1)} - u^{(2)}$ ,  $[u]|_{S_3} = u^{(3)} - u^{(2)}$ ,  $u^{(i)} = u(x, t)|_{x \in \bar{\Omega}_i}$ .

By  $\mathbf{n}_x = (\mathbf{n}_1, \mathbf{n}_2, \mathbf{n}_3)$  and  $\mathbf{n}_t$  we mean the components of the normal vector  $\mathbf{n}$  to the surface  $\mathfrak{G} = \{x \in \Gamma_t, t > 0\}$  in  $\mathbb{R}^4$ .

We assume that  $\Gamma_0$  is located in the neighborhood of a smooth connected surface  $\mathcal{G}$  of arbitrary shape and can be regarded as a normal perturbation of  $\mathcal{G}$ :

$$\Gamma_0 = \{x = y + \mathbf{N}(y)\rho_0(y), \quad y \in \mathcal{G}\},$$

where  $\rho_0$  is a given small function and  $\mathbf{N}(y)$  is the normal to  $\mathcal{G}$  exterior with respect to  $\Omega_0$ . Moreover, we assume that also for  $t > 0$

$$\Gamma_t = \{x = y + \mathbf{N}(y)\rho(y, t), \quad y \in \mathcal{G}\},$$

with an unknown function  $\rho(y, t)$  such that  $\rho(y, 0) = \rho_0(y)$ . We extend  $\mathbf{N}(y)$  and  $\rho(y, t)$  from  $\mathcal{G}$  into  $\Omega$  in such a way that the extension  $\mathbf{N}^*$  of  $\mathbf{N}$  is a smooth non-zero regular function in  $\Omega$  and  $\rho^*$  vanishes near  $S \cup S_3$  and satisfies the inequalities (1.23) (hence  $\rho^*$  is small for small  $\rho$ ).

Let  $\mathcal{F}_1$  be the domain bounded by  $\mathcal{G}$ ,  $\mathcal{F}_3 = \Omega_3$ ,  $\mathcal{F}_2 = \Omega \setminus (\bar{\mathcal{F}}_1 \cup \bar{\mathcal{F}}_3)$ . The transformation

$$x \equiv e_\rho(y, t) = y + \mathbf{N}^*(y)\rho^*(y, t) \quad (1.5)$$

maps  $\mathcal{F}_1$  on  $\Omega_{1t}$ ,  $\mathcal{F}_2$  on  $\Omega_{2t}$ ,  $\mathcal{F}_3$  on itself and, as shown in [2], it converts (1.2)-(1.4) in

$$\begin{cases} \mathbf{u}_t - \rho_t^*(\mathcal{L}^{-1}\mathbf{N}^*(y) \cdot \nabla)\mathbf{u} + (\mathcal{L}^{-1}\mathbf{u} \cdot \nabla)\mathbf{u} \\ - \tilde{\nabla} \cdot \tilde{T}(\mathbf{u}, q) - \tilde{\nabla} \cdot T_M(\frac{\mathcal{L}}{L}\mathbf{h}) = \mathbf{f}(e_\rho, t), \\ \nabla \cdot \hat{\mathcal{L}}\mathbf{u} = 0, \quad y \in \mathcal{F}_1, \quad t > 0, \\ \tilde{T}(\mathbf{u}, q)\mathbf{n}(e_\rho, t) + [T(\frac{\mathcal{L}}{L}\mathbf{h})\mathbf{n}(e_\rho, t)] = \sigma H\mathbf{n}, \quad \rho_t = \frac{\mathbf{u} \cdot \hat{\mathcal{L}}^T \mathbf{N}}{\Lambda(y, \rho)}, \quad y \in \mathcal{G}, \\ \mathbf{u}(y, 0) = \mathbf{u}_0(y) = \mathbf{v}_0(e_{\rho_0}), \quad y \in \mathcal{F}_1, \quad \rho(y, 0) = \rho_0(y), \quad y \in \mathcal{G}, \end{cases} \quad (1.6)$$

$$\begin{cases} \mu(\mathbf{h}_t - \frac{1}{L}\hat{\mathcal{L}}_t\mathcal{L}\mathbf{h} - \rho_t^*\hat{\mathcal{L}}(\mathcal{L}^{-1}\mathbf{N}^*(y) \cdot \nabla)\frac{1}{L}\mathcal{L}\mathbf{h}) = -\text{rot}\mathcal{P}\mathbf{e}, \quad y \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3, \\ \mathcal{P}\text{rot}\mathcal{P}\mathbf{h} = \alpha(\mathcal{P}\mathbf{e} + \mu(\mathcal{L}^{-1}\mathbf{u} \times \mathbf{h})), \quad \nabla \cdot \mathbf{h} = 0, \quad y \in \mathcal{F}_1, \\ \text{rot}\mathcal{P}\mathbf{h} = 0, \quad \nabla \cdot \mathbf{h} = 0, \quad \nabla \cdot \mathbf{e} = 0, \quad x \in \mathcal{F}_2, \\ \text{roth} = \alpha\mathbf{e} + \mathbf{j}(y, t), \quad \nabla \cdot \mathbf{h} = 0, \quad y \in \mathcal{F}_3, \end{cases} \quad (1.7)$$

$$\begin{cases} \mathbf{h} \cdot \mathbf{n} = 0, \quad \mathbf{e}_\tau = 0, \quad y \in S, \\ [\mu\mathbf{h} \cdot \mathbf{n}] = 0, \quad [\mathbf{h}_\tau] = 0, \quad [\mathbf{e}_\tau] = 0, \quad y \in S_3, \\ [\mu\mathbf{h} \cdot \mathbf{N}] = 0, \quad [\mathbf{h} - \frac{\tilde{\mathcal{L}}\tilde{\mathcal{L}}^T \mathbf{N}}{|\tilde{\mathcal{L}}^T \mathbf{N}|^2}(\mathbf{h} \cdot \mathbf{N})] = 0, \\ -\Lambda(y, \rho)\rho_t[\mu\mathbf{h}] + L[\mathbf{N} \times \mathcal{P}\mathbf{e}] = 0, \quad y \in \mathcal{G}, \\ \mathbf{h}_0(y, 0) = \mathbf{h}_0(y) = \hat{\mathcal{L}}(y, \rho_0^*)\mathbf{H}(e_{\rho_0}), \quad y \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3, \end{cases} \quad (1.8)$$

where

$$\mathbf{u}(y, t) = \mathbf{v}(e_\rho, t), \quad q(y, t) = p(e_\rho, t), \quad \mathbf{h} = \hat{\mathcal{L}}\mathbf{H}(e_\rho, t), \quad \mathbf{e} = \hat{\mathcal{L}}\mathbf{E}(e_\rho, t), \quad (1.9)$$

$\mathcal{L} = \mathcal{L}(y, \rho^*) = (l_{ij})_{i,j=1,2,3}$  is the Jacobi matrix of the transformation (1.5),  $L = \det \mathcal{L}$ ,

$\hat{\mathcal{L}} = L\mathcal{L}^{-1}$  is the co-factors matrix of  $\mathcal{L}$ ;

$\mathcal{P}(y, \rho^*) = \frac{\mathcal{L}^T}{L}\mathcal{L}$ ,

$\tilde{\nabla} = \mathcal{L}^{-T}\nabla_y$  is the transformed gradient  $\nabla_x$  ("T" means transposition,  $\mathcal{L}^{-T} = (\mathcal{L}^{-1})^T$ );

$\tilde{S}(\mathbf{u}) = \tilde{\nabla}\mathbf{u} + (\tilde{\nabla}\mathbf{u})^T$  is the transformed rate-of-strain tensor,

$\tilde{T}(\mathbf{u}, q) = -q\mathbf{I} + \nu\tilde{S}(\mathbf{u})$  is the transformed stress tensor,

$\Lambda(y, \rho) = \mathbf{N}(y) \cdot \hat{\mathcal{L}}(y, \rho)\mathbf{N}(y) = 1 - \rho\mathcal{H}(y) + \rho^2\mathcal{K}(y)$

and  $\mathcal{H}, \mathcal{K}$  are the doubled mean curvature and the Gaussian curvature of  $\mathcal{G}$ , respectively.

We study the problem (1.6)-(1.8) in the spaces

$$W_p^{2,1}(D_T) = W_p^{2,0}(D_T) \cap W_p^{0,1}(D_T), \quad p > 3, \quad D_T = D \times (0, T), \quad D \subset \mathbb{R}^n,$$

where  $W_p^{2,0}(D_T) = L_p(0, T; W_p^2(D))$ ,  $W_p^{0,1}(D_T) = W_p^1(0, T; L_p(D))$ . The norm in  $W_p^{2,1}(D_T)$  is defined by

$$\|u\|_{W_p^{2,1}(D_T)}^p = \int_0^T \|u(\cdot, t)\|_{W_p^2(D)}^p dt + \int_0^T (\|u_t(\cdot, t)\|_{L_p(D)}^p + \|u(\cdot, t)\|_{L_p(D)}^p) dt, \quad (1.10)$$

and the integrals in the right-hand side represent the norms in  $W_p^{2,0}(D_T)$  and  $W_p^{0,1}(D_T)$ , respectively.

By  $W_p^l(D)$  we mean the space of functions  $u(x)$ ,  $x \in D$ , with finite norm

$$\|u\|_{W_p^l(D)} = \left( \sum_{|j| \leq l} \|D^j u\|_{L_p(D)}^p \right)^{1/p},$$

if  $l$  is an integer, and

$$\|u\|_{W_p^l(D)} = \left( \|u\|_{W_p^{[l]}(D)}^p + \sum_{|j|=l} \int_D \int_D \frac{|D^j u(x) - D^j u(y)|^p}{|x - y|^{n+p\lambda}} dx dy \right)^{1/p},$$

if  $l = [l] + \lambda$ ,  $0 < \lambda < 1$ . The anisotropic space  $W_p^{l,l/2}(D_T)$  is defined as  $W_p^{l,0}(D_T) \cap W_p^{0,1/2}(D_T) = L_p(0, T; W_p^l(D)) \cap W_p^{l/2}(0, T; L_p(D))$ .

The spaces  $W_p^l$ ,  $W_p^{l,l/2}$  on smooth manifolds are defined, as usual, with the help of local maps and partition of unity.

We recall the trace and extension theorems for the space  $W_p^{2,1}(D_T)$  (see [6,7]): if  $u \in W_p^{2,1}(D_T)$ , then  $u|_{t=t_0} \in W_p^{2-2/p}(D)$ ,  $u|_{x \in \partial D} \in W_p^{2-1/p, 1-1/(2p)}(\partial D \times (0, T))$ , and

$$\|u(\cdot, t_0)\|_{W_p^{2-2/p}(D)} \leq c \|u\|_{W_p^{2,1}(D_T)},$$

$$\|u\|_{W_p^{2-1/p, 1-1/(2p)}(\partial D \times (0, T))} \leq c \|u\|_{W_p^{2,1}(D_T)}.$$

For arbitrary  $\varphi \in W_p^{2-2/p}(D)$  there exists such  $u \in W_p^{2,1}(D_T)$  that  $u(x, 0) = \varphi(x)$  and

$$\|u\|_{W_p^{2,1}(D_T)} \leq c \|\varphi\|_{W_p^{2-2/p}(D)};$$

for arbitrary  $\psi \in W_p^{2-1/p, 1-1/(2p)}(\partial D \times (0, T))$  there exists such  $u \in W_p^{2,1}(D_T)$  that  $u|_{x \in \partial D} = \psi(x, t)$  and

$$\|u\|_{W_p^{2,1}(D_T)} \leq c \|\psi\|_{W_p^{2-1/p, 1-1/(2p)}(\partial D \times (0, T))}.$$

Moreover, if  $\rho \in W_p^{3-1/p, 0}(\partial D \times (0, T)) \cap W_p^1(0, T; W_p^{2-1/p}(\partial D))$ , then  $\rho(\cdot, t_0) \in W_p^{3-2/p}(\partial D)$  and

$$\|\rho(\cdot, t_0)\|_{W_p^{3-2/p}(\partial D)} \leq c (\|\rho\|_{W_p^{3-1/p, 0}(\partial D \times (0, T))} + \|\rho_t\|_{W_p^{2-1/p, 0}(\partial D \times (0, T))}). \quad (1.11)$$

For arbitrary  $\rho_0 \in W_p^{3-2/p}(\partial D)$  there exists  $\rho \in W_p^{3-1/p, 0}(\partial D \times (0, T))$  with  $\rho_t \in W_p^{2-1/p, 1-1/2p}(\partial D \times (0, T))$  such that  $\rho(x, 0) = \rho_0(x)$  and

$$\|\rho\|_{W_p^{3-1/p, 0}(\partial D \times (0, T))} + \|\rho_t\|_{W_p^{2-1/p, 1-1/2p}(\partial D \times (0, T))} \leq c \|\rho_0\|_{W_p^{3-2/p}(\partial D)}.$$

Following [8,9], we introduce some spaces of divergence free vector fields. Let  $\mathcal{D}$  be a bounded domain in  $\mathbb{R}^3$  with a smooth boundary. We define finite-dimensional spaces

$$\begin{aligned} U_n(\mathcal{D}) &= \{\mathbf{u} \in W_p^1(\mathcal{D}), \quad \nabla \cdot \mathbf{u} = 0, \quad \text{rot} \mathbf{u} = 0, \quad \mathbf{u} \cdot \mathbf{n}|_{\partial \mathcal{D}} = 0\}, \\ U_d(\mathcal{D}) &= \{\mathbf{u} \in W_p^1(\mathcal{D}), \quad \nabla \cdot \mathbf{u} = 0, \quad \text{rot} \mathbf{u} = 0, \quad \mathbf{u}_\tau|_{\partial \mathcal{D}} = 0\} \end{aligned}$$

of the Neumann and Dirichlet vector fields. The dimensions of these spaces are equal to the first and the second Betti number of the domain  $\mathcal{D}$ ,  $b_1(\mathcal{D})$  and  $b_2(\mathcal{D})$ , respectively.

If  $b_1(\mathcal{D}) > 0$ , then there exist  $b_1(\mathcal{D})$  smooth closed contours in  $\mathcal{D}^c = \mathbb{R}^3 \setminus \overline{\mathcal{D}}$ , generating the first homology group of  $\mathcal{D}^c$ , and every such contour  $\Lambda$  generates the Neumann vector field of the form

$$\mathbf{u}(x) = \mathbf{u}_1(x) + \mathbf{u}_2(x), \quad x \in \mathcal{D}, \quad (1.12)$$

where  $\mathbf{u}_1$  is defined through the Bio-Savard law:

$$\mathbf{u}_1(x) = \int_{\Lambda} \frac{x - y}{|x - y|^3} \times d\mathbf{l}_y,$$

and  $\mathbf{u}_2 = \nabla \varphi$ ,

$$\nabla^2 \varphi(x) = 0, \quad x \in \mathcal{D}, \quad \frac{\partial \varphi}{\partial n}|_{\partial \mathcal{D}} = -\mathbf{u}_1 \cdot \mathbf{n}|_{\partial \mathcal{D}}.$$

As for  $b_2(\mathcal{D})$ , this number is equal to the number of the connected components of the boundary  $\partial \mathcal{D}$  of  $\mathcal{D}$  minus 1, and if  $\partial \mathcal{D} = \bigcup_{k=0}^{b_2(\mathcal{D})} \Sigma_k$ , then the basis in  $U_d(\mathcal{D})$  is formed by the vector fields  $\mathbf{v}_k = \nabla \Phi_k$ , where  $\Phi_k$  are the solutions to the problems

$$\nabla^2 \Phi_k = 0, \quad \Phi_k|_{\Sigma_j} = \delta_{jk}, \quad k, j = 1, \dots, b_2(\mathcal{D}), \quad \Phi_k|_{\Sigma_0} = 0. \quad (1.13)$$

If  $\Sigma_k \in C^m$ , then  $\mathbf{v}_k, \mathbf{u}_j \in W_p^{m-1}(\mathcal{D})$  for arbitrary  $p > 1$ .

For the domain  $\Omega = \bar{\mathcal{F}}_1 \cup \mathcal{F}_2 \cup \bar{\mathcal{F}}_3$  introduced above we have

$$b_1(\mathcal{F}_2) = b_1(\Omega) + b_1(\mathcal{F}_1 \cup \mathcal{F}_3).$$

The elements  $\mathbf{u}_j \in U_n(\mathcal{F}_2)$  can be generated by the contours  $\Lambda_j \subset \mathbb{R}^3 \setminus \bar{\Omega}$  (in this case  $j = 1, \dots, b_1(\Omega)$ ) or  $\Lambda_j \subset \mathcal{F}_1, \Lambda_j \subset \mathcal{F}_3$  ( $j = b_1(\Omega) + 1, \dots, b_1(\Omega) + b_1(\mathcal{F}_1 \cup \mathcal{F}_3)$ ). It is easily seen that in the first case  $\mathbf{u}_j^{\mathcal{F}} \in U_n(\mathcal{F}_2)$  are connected with  $\mathbf{u}_j^{\Omega} \in U_n(\Omega)$  by

$$\begin{aligned} \mathbf{u}_j^{\mathcal{F}}(x) &= \mathbf{u}_j^{\Omega}(x) + \nabla \omega_j(x), \quad x \in \mathcal{F}_2, \\ \nabla^2 \omega_j(x) &= 0, \quad x \in \mathcal{F}_2, \quad \frac{\partial \omega_j(x)}{\partial n} = 0, \quad x \in S, \\ \frac{\partial \omega_j(x)}{\partial N} &= -\mathbf{u}_j^{\Omega} \cdot \mathbf{N}, \quad x \in \mathcal{G}, \quad \frac{\partial \omega_j(x)}{\partial n} = -\mathbf{u}_j^{\Omega} \cdot \mathbf{n}, \quad x \in S_3. \end{aligned}$$

Finally, we introduce in  $\Omega = \bar{\mathcal{F}}_1 \cup \mathcal{F}_2 \cup \bar{\mathcal{F}}_3$  the space  $\tilde{U}_n(\Omega)$  of the modified Neumann vector fields  $\tilde{\mathbf{u}}_q, q = 1, \dots, b_1(\Omega)$ , satisfying the conditions

$$\left\{ \begin{aligned} &\text{rot} \tilde{\mathbf{u}}_q(x) = 0, \quad \nabla \cdot \tilde{\mathbf{u}}_q(x) = 0, \quad x \in \mathcal{F}_i, \quad i = 1, 2, 3, \\ &[\mu \tilde{\mathbf{u}}_q \cdot \mathbf{N}] = 0, \quad [\tilde{\mathbf{u}}_{q,\tau}] = 0, \quad x \in \mathcal{G}, \quad [\mu \tilde{\mathbf{u}}_q \cdot \mathbf{n}] = 0, \quad [\tilde{\mathbf{u}}_{q,\tau}] = 0, \quad x \in S_3, \\ &\tilde{\mathbf{u}}_q \cdot \mathbf{n} = 0, \quad x \in S. \end{aligned} \right. \quad (1.14)$$

It is clear that

$$\begin{aligned}\tilde{\mathbf{u}}_q &= \mathbf{u}_q + \nabla \phi_q(x), \quad \mathbf{u}_q \in U_n(\Omega), \\ \nabla^2 \phi_q(x) &= 0, \quad x \in \mathcal{F}_i, \quad i = 1, 2, 3, \quad \frac{\partial \phi_q}{\partial n} = 0, \quad x \in S, \\ [\phi_q] &= 0, \quad [\mu \frac{\partial \phi_q}{\partial N}] = -[\mu] \mathbf{u}_q \cdot \mathbf{N}, \quad x \in \mathcal{G}, \quad [\phi_q] = 0, \quad [\mu \frac{\partial \phi_q}{\partial n}] = -[\mu] \mathbf{u}_q \cdot \mathbf{n}, \quad x \in S_3.\end{aligned}$$

Let us go back to the problem (1.6)-(1.8). We separate the determination of  $(\mathbf{u}, q, \rho, \mathbf{h})$  from that of  $\mathbf{e}$ . It is easily seen that (1.7), (1.8) imply

$$\begin{cases} \mu(\mathbf{h}_t - \Phi) + \alpha^{-1} \text{rot} \mathcal{P} \text{rot} \mathcal{P} \mathbf{h} = \text{rot} \mathbf{J}, & \nabla \cdot \mathbf{h}(y, t) = 0, \quad y \in \mathcal{F}_1 \cup \mathcal{F}_3, \\ \text{rot} \mathcal{P} \mathbf{h}(y, t) = 0, & \nabla \cdot \mathbf{h} = 0, \quad y \in \mathcal{F}_2, \\ [\mu \mathbf{h} \cdot \mathbf{N}] = 0, \quad [\mathbf{h}_\tau] = \left( \frac{\widehat{\mathcal{L}} \widehat{\mathcal{L}}^T \mathbf{N}}{|\widehat{\mathcal{L}}^T \mathbf{N}|^2} - \mathbf{N} \right) [\mathbf{h} \cdot \mathbf{N}], & y \in \mathcal{G}, \\ [\mu \mathbf{h} \cdot \mathbf{n}] = 0, \quad [\mathbf{h}_\tau] = 0, & y \in S_3, \quad \mathbf{h} \cdot \mathbf{n} = 0, \quad y \in S, \\ \mathbf{h}(y, 0) = \mathbf{h}_0(y), & y \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3, \end{cases} \quad (1.15)$$

where  $\mathbf{J} = \mu_1 \mathcal{L}^{-1} \mathbf{u} \times \mathbf{h}$  in  $\mathcal{F}_1$ ,  $\mathbf{J} = \alpha^{-1} \mathbf{j}(y, t)$  in  $\mathcal{F}_3$ , and

$$\Phi = \frac{1}{L} \widehat{\mathcal{L}}_t \mathcal{L} \mathbf{h} + \rho_t^* \widehat{\mathcal{L}} (\mathcal{L}^{-1} \mathbf{N}^*(y) \cdot \nabla) \frac{1}{L} \mathcal{L} \mathbf{h}.$$

As shown in [2], the vector field  $\Phi$  is divergence free.

In the case  $b_1(\mathcal{F}_2) > 0$  equations (1.15) should be supplemented by some orthogonality conditions. Multiplying the first equation in (1.7) by  $\mathbf{u}_q^\mathcal{F} \in U_n(\mathcal{F}_2)$  and integrating over  $\mathcal{F}_2$  we obtain

$$\begin{aligned} \int_{\mathcal{F}_2} \mu(\mathbf{h}_t - \Phi) \cdot \mathbf{u}_q^\mathcal{F} dy &= - \int_{\mathcal{F}_2} \text{rot} \mathcal{P} \mathbf{e}^{(2)} \cdot \mathbf{u}_q^\mathcal{F} dy = \int_{\mathcal{G}} (\mathbf{N} \times \mathcal{P} \mathbf{e}^{(2)}) \cdot \mathbf{u}_q^\mathcal{F} dS \\ &+ \int_{S_3} (\mathbf{n} \times \mathbf{e}^{(2)}) \cdot \mathbf{u}_q^\mathcal{F} dS = \int_{\mathcal{G}} (\mathbf{N} \times \mathcal{P} \mathbf{e}^{(1)}) \cdot \mathbf{u}_q^\mathcal{F} dS + \int_{S_3} (\mathbf{n} \times \mathbf{e}^{(3)}) \cdot \mathbf{u}_q^\mathcal{F} dS - \int_{\mathcal{G}} \Psi \cdot \mathbf{u}_q^\mathcal{F} dS \\ &= \int_{\mathcal{G}} (\mathbf{N} \times (\alpha^{-1} \mathcal{P} \text{rot} \mathcal{P} \mathbf{h}^{(1)} - \mathbf{J})) \cdot \mathbf{u}_q^\mathcal{F} dS + \int_{S_3} (\mathbf{n} \times \alpha^{-1} (\text{rot} \mathbf{h}^{(3)} - \mathbf{j}(y, t))) \cdot \mathbf{u}_q^\mathcal{F}(y, t) dS \\ &- \int_{\mathcal{G}} \Psi \cdot \mathbf{u}_q^\mathcal{F} dS, \quad q = 1, \dots, b_1(\mathcal{F}_2), \end{aligned} \quad (1.16)$$

where  $\mathbf{h}^{(i)} = \mathbf{h}|_{y \in \overline{\mathcal{F}}_i}$ ,  $\mathbf{e}^{(i)} = \mathbf{e}|_{y \in \overline{\mathcal{F}}_i}$ ,

$$\Psi = \frac{\rho_t \Lambda(y, \rho)}{L(y, \rho)} [\mu \mathbf{h}], \quad y \in \mathcal{G}.$$

The same equations are satisfied for  $\mathbf{u}_q^\Omega(x)$ ,  $q = 1, \dots, b_1(\Omega)$ :

$$\begin{aligned} \int_{\mathcal{F}_2} \mu(\mathbf{h}_t - \Phi) \cdot \mathbf{u}_q^\Omega dy &= \int_{\mathcal{G}} \mathbf{N} \times (\alpha^{-1} \mathcal{P} \text{rot} \mathcal{P} \mathbf{h}^{(1)} - \mathbf{J}) \cdot \mathbf{u}_q^\Omega dS \\ &+ \int_{S_3} \mathbf{n} \times \alpha^{-1} (\text{rot} \mathbf{h}^{(3)} - \mathbf{j}(y, t)) \cdot \mathbf{u}_q^\Omega(y, t) dS - \int_{\mathcal{G}} \Psi \cdot \mathbf{u}_q^\Omega dS \end{aligned} \quad (1.17)$$

and, since

$$\int_{\mathcal{F}_1 \cup \mathcal{F}_3} \mu(\mathbf{h}_t - \Phi) \cdot \mathbf{u}_q^\Omega dy = \int_{\mathcal{F}_1 \cup \mathcal{F}_3} (-\alpha^{-1} \text{rot} \mathcal{P} \text{rot} \mathcal{P} \mathbf{h} + \text{rot} \mathbf{J}) \cdot \mathbf{u}_q^\Omega dy$$



$$= - \int_{\mathcal{G}} \mathbf{N} \times (\alpha^{-1} \mathcal{P} \text{rot} \mathcal{P} \mathbf{h}^{(1)} - \mathbf{J}) \cdot \mathbf{u}_q^\Omega dS - \int_{S_3} \mathbf{n} \times \alpha^{-1} (\text{rot} \mathbf{h}^{(3)} - \mathbf{j}) \cdot \mathbf{u}_q^\Omega dS,$$

it holds

$$\int_{\Omega} \mu(\mathbf{h}_t - \Phi) \cdot \mathbf{u}_q^\Omega dy = - \int_{\mathcal{G}} \Psi \cdot \mathbf{u}_q^\Omega dS, \quad q = 1, \dots, b_1(\Omega). \quad (1.18)$$

Hence, in addition to (1.15), we have (1.16), (1.18).

Equations (1.6), (1.15), (1.16) constitute the main problem for  $\mathbf{u}, q, \rho, \mathbf{h}$  that is solved in Sec. 2 and 3. Making use of (1.18), we construct in Sec.4  $\mathbf{e}(y, t)$  satisfying (1.7), (1.8).

**Theorem 1.** *Let  $\mathbf{u}_0 \in W_p^{2-2/p}(\mathcal{F}_1)$ ,  $\rho_0 \in W_p^{3-2/p}(\mathcal{G})$ ,  $\mathbf{h}_0 \in W_r^{2-2/r}(\mathcal{F}_i)$ ,  $i = 1, 2, 3$ , with*

$$3 < r < p, \quad 1/r - 1/p \leq 1/5(1 - 2/p), \quad (1.19)$$

*and let the compatibility conditions*

$$\begin{aligned} \tilde{\nabla} \cdot \mathbf{u}_0 &= 0, \quad y \in \mathcal{F}_1, \quad \tilde{S}(\mathbf{u}_0) \mathbf{n}_0(e_{\rho_0}) - \mathbf{n}_0(\mathbf{n}_0 \cdot \tilde{S}(\mathbf{u}_0) \mathbf{n}_0) = 0, \quad y \in \mathcal{G}, \\ \nabla \cdot \mathbf{h}_0 &= 0, \quad y \in \mathcal{F}_i, \quad i = 1, 2, 3, \\ \text{rot} \mathcal{P}(y, \rho_0^*) \mathbf{h}_0 &= 0, \quad y \in \mathcal{F}_2, \\ [\mu \mathbf{h}_0 \cdot \mathbf{N}] &= 0, \quad [\mathbf{h}_{0\tau}] = \left( \frac{\hat{\mathcal{L}}(y, \rho_0) \hat{\mathcal{L}}^T \mathbf{N}}{|\hat{\mathcal{L}}^T \mathbf{N}|^2} - \mathbf{N} \right) [\mathbf{h}_0 \cdot \mathbf{N}], \quad y \in \mathcal{G}, \\ [\mu \mathbf{h}_0 \cdot \mathbf{n}] &= 0, \quad [\mathbf{h}_{0\tau}] = 0, \quad y \in S_3, \quad \mathbf{h}_0 \cdot \mathbf{n} = 0, \quad y \in S, \end{aligned} \quad (1.20)$$

*where  $\mathbf{n}_0$  is the normal to  $\Gamma_0$ , as well as the smallness condition*

$$\|\rho_0\|_{W_p^{2-1/p}(\mathcal{G})} \leq \epsilon \ll 1 \quad (1.21)$$

*be satisfied. Assume also that*

$$\mathbf{f} \in W_p^1(\mathbb{R}^3), \quad \forall t \in (0, T_0), \quad \mathbf{j} \in W_r^{1,0}(\mathcal{F}_3 \times (0, T_0)), \quad (1.22)$$

*(1.1) holds and the extension  $\rho^*$  of  $\rho$  in (1.5) satisfies the conditions*

$$\begin{aligned} \frac{\partial \rho^*(y, t)}{\partial N} \Big|_{\mathcal{G}} &= 0, \quad \rho^*(y, t) = 0 \text{ near } S \cup S_3 \text{ and in } \mathcal{F}_3, \\ \|\rho^*(\cdot, t)\|_{W_p^l(\Omega)} &\leq c \|\rho\|_{W_p^{l-1/p}(\mathcal{G})}, \quad l \in (1/p, 3], \\ \|\rho_t^*(\cdot, t)\|_{W_p^l(\Omega)} &\leq c \|\rho_t\|_{W_p^{l-1/p}(\mathcal{G})}, \quad r \in (1/p, 2]. \end{aligned} \quad (1.23)$$

*Then the problem (1.6), (1.15), (1.16) has a unique solution defined in a certain (small) time interval  $(0, T)$ ,  $T \leq T_0$ , with the following regularity properties:*

$$\begin{aligned} \mathbf{u} &\in W_p^{2,1}(Q_T^1), \quad \nabla q \in L_p(Q_T^1), \quad q \in W_p^{l-1/p,0}(G_T), \quad \rho \in W_p^{3-3/p,0}(G_T), \\ \rho_t &\in W_p^{2-1/p,1-1/2p}(G_T), \quad \mathbf{h} \in W_r^{2,1}(Q_T^i), \quad i = 1, 2, 3, \end{aligned}$$

where  $Q_T^i = \mathcal{F}_i \times (0, T)$ ,  $G_T = \mathcal{G} \times (0, T)$ . The solution satisfies the inequality

$$\begin{aligned}
& \|\mathbf{u}\|_{W_p^{2,1}(Q_T^1)} + \|\nabla q\|_{L_p(Q_T^1)} + \|q\|_{W_p^{1-1/p,0}(G_T)} + \sup_{t < T} \|\mathbf{u}(\cdot, t)\|_{W_p^{2-2/p}(\mathcal{F}_1)} + \sup_{t < T} \|\rho\|_{W_p^{3-2/p}(\mathcal{G})} \\
& + \|\rho\|_{W_p^{3-1/p,0}(G_T)} + \|\rho_t\|_{W_p^{2-1/p,1-1/2p}(G_T)} + \sum_{i=1}^3 \|\mathbf{h}^{(i)}\|_{W_r^{2,1}(Q_T^i)} \\
& + \sum_{i=1}^3 \sup_{t < T} \|\mathbf{h}\|_{W_r^{2-2/r}(\mathcal{F}_i)} \leq c \left( \|\mathbf{f}\|_{L_p(Q_T^1)} + \|\mathcal{H}\|_{W_p^{1-1/p}(\mathcal{G})} + \|\mathbf{u}_0\|_{W_p^{2-2/p}(\mathcal{F}_1)} \right. \\
& \left. + \|\rho_0\|_{W_p^{3-2/p}(\mathcal{G})} + \sum_{i=1}^3 \|\mathbf{h}_0\|_{W_r^{2-2/r}(\Omega_{i0})} + \|\mathbf{j}\|_{W_r^{1,0}(Q_T^3)} \right). \tag{1.24}
\end{aligned}$$

Once the solution with the above-mentioned properties is obtained, it can be shown that

$$\mathbf{h}^{(i)} \in W_p^{2,1}(Q_T^i), \quad i = 1, 2, 3,$$

provided  $\mathbf{h}_0 \in W_p^{2-2/p}(\Omega_{i0})$ ,  $i = 1, 2, 3$ ,  $\mathbf{j} \in W_p^{1,0}(Q_T^3)$  (cf. [3, Sec. 5]).

## 2 Linear problems

The proof of Theorem 1 is based on the analysis of the following non-homogeneous linear problems:

1. Find  $(\mathbf{v}, p, \rho)$  such that

$$\begin{cases} \mathbf{v}_t - \nu \nabla^2 \mathbf{v} + \nabla p = \mathbf{f}(y, t), \\ \nabla \cdot \mathbf{v} = f(y, t) = \nabla \cdot \mathbf{F}(y, t), \quad y \in \mathcal{F}_1, \quad t > 0, \\ T(\mathbf{v}, p) \mathbf{N}(y) + \sigma \mathbf{N}(y) \mathfrak{B} \rho = \mathbf{d}(y, t), \\ \rho_t + \mathbf{V}(x) \cdot \nabla_\tau \rho - \mathbf{v} \cdot \mathbf{N}(y) = g(y, t), \quad y \in \mathcal{G}, \\ \mathbf{v}(y, 0) = \mathbf{v}_0(y), \quad y \in \mathcal{F}_1, \quad \rho(y, 0) = \rho_0(y), \quad y \in \mathcal{G}, \end{cases} \tag{2.1}$$

where  $\mathfrak{B} \rho = -\Delta_{\mathcal{G}} \rho - b(y) \rho$ ,  $b = (\mathcal{H}^2 - 2\mathcal{K})$ ,  $\Delta_{\mathcal{G}}$  is the Laplace-Beltrami operator on  $\mathcal{G}$ ,  $\mathbf{V}$  is a given vector field from  $W_p^{2-1/p}(\mathcal{G})$ .

2. Find the vector field  $\mathbf{H}(y, t)$ , satisfying the equations

$$\begin{cases} \mu \mathbf{H}_t(y, t) + \alpha^{-1} \text{rot} \text{rot} \mathbf{H}(y, t) = \mathbf{G}(y, t), \quad \nabla \cdot \mathbf{H}(y, t) = 0, \quad y \in \mathcal{F}_1 \cup \mathcal{F}_3, \\ \text{rot} \mathbf{H}(y, t) = \text{rot} \ell(y, t), \quad \nabla \cdot \mathbf{H}(y, t) = 0, \quad y \in \mathcal{F}_2 \\ [\mu \mathbf{H} \cdot \mathbf{N}] = 0, \quad [\mathbf{H}_\tau] = \mathbf{a}(y, t), \quad y \in \mathcal{G}, \\ [\mu \mathbf{H} \cdot \mathbf{n}] = 0, \quad [\mathbf{H}_\tau] = 0, \quad y \in S_3, \quad \mathbf{H} \cdot \mathbf{n} = 0, \quad y \in S, \\ \int_{\mathcal{F}_2} \mu \mathbf{H}_t \cdot \mathbf{u}_q^{\mathcal{F}} dy = \int_{\mathcal{G}} (\mathbf{N} \times \alpha^{-1} \text{rot} \mathbf{H}^{(1)}) \cdot \mathbf{u}_q^{\mathcal{F}}(y) dS \\ + \int_{S_3} (\mathbf{n} \times \alpha^{-1} \text{rot} \mathbf{H}^{(3)}) \cdot \mathbf{u}_q^{\mathcal{F}}(y) dS + \mathbf{M}_q(t), \quad q = 1, \dots, b_1(\mathcal{F}_2), \\ \mathbf{H}(y, 0) = \mathbf{H}_0(y), \quad y \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3. \end{cases} \tag{2.2}$$

In addition, we need to consider the auxiliary problem

3.

$$\begin{cases} \operatorname{rot} \mathbf{h}(y) = \mathbf{k}(y), & \nabla \cdot \mathbf{h} = 0, & y \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3, \\ [\mu \mathbf{h} \cdot \mathbf{N}] = 0, & [\mathbf{h}_\tau] = \mathbf{a}, & y \in \mathcal{G}, \\ [\mu \mathbf{h} \cdot \mathbf{n}] = 0, & [\mathbf{h}_\tau] = 0, & y \in S_3, \\ \mathbf{h} \cdot \mathbf{n}(y) = 0, & & y \in S. \end{cases} \quad (2.3)$$

**Theorem 2.** Assume that

$$\begin{aligned} \mathbf{f} &\in L_p(Q_T^1), \quad f \in W_p^{1,0}(Q_T^1), \quad f = \nabla \mathbf{F}, \quad \mathbf{F} \in W_p^{0,1}(Q_T^1), \\ \mathbf{d} \cdot \mathbf{N} &\in W_2^{1-1/p,0}(G_T), \quad \mathbf{d} - \mathbf{N}(\mathbf{d} \cdot \mathbf{N}) \equiv \mathbf{d}_\tau \in W_p^{1-1/p,1/2-1/(2p)}(G_T), \\ \mathbf{g} &\in W_p^{2-1/p,1-1/2p}(G_T), \quad \mathbf{v}_0 \in W_p^{2-2/p}(\mathcal{F}_1), \\ \rho_0 &\in W_p^{3-2/p}(\mathcal{G}), \quad \mathbf{V} \in W_p^{2-1/p}(\mathcal{G}) \end{aligned}$$

with  $p > 3$  and let the compatibility conditions

$$\nabla \cdot \mathbf{v}_0(x) = f(x, 0), \quad x \in \mathcal{F}_1, \quad \nu(S(\mathbf{v}_0)\mathbf{N})_\tau = \mathbf{d}_\tau(x, 0), \quad x \in \mathcal{G}, \quad (2.4)$$

be satisfied. Then the problem (2.1) has a unique solution  $\mathbf{v}, p, \rho$  such that  $\mathbf{v} \in W_p^{2,1}(Q_T^1)$ ,  $\nabla p \in L_p(Q_T^1)$ ,  $p \in W_p^{1-1/p,0}(G_T)$ ,  $\rho \in W_p^{3-1/p,0}(G_T)$ ,  $\rho_t \in W_p^{2-1/p,1-1/2p}(G_T)$ ,  $\rho(\cdot, t) \in W_p^{3-2/p}(\mathcal{G})$ ,  $\forall t \in (0, T)$ , and the solution satisfies the inequality

$$\begin{aligned} &\|\mathbf{v}\|_{W_p^{2,1}(Q_T^1)} + \|\nabla p\|_{L_p(Q_T^1)} + \|p\|_{W_p^{1-1/p,0}(G_T)} + \sup_{t < T} \|\mathbf{v}(\cdot, t)\|_{W_p^{2-2/p}(\mathcal{F}_1)} \\ &+ \|\rho\|_{W_p^{3-1/p,0}(G_T)} + \|\rho_t\|_{W_2^{2-1/p,1-1/2p}(G_T)} + \sup_{t < T} \|\rho\|_{W_p^{3-2/p}(\mathcal{G})} \\ &\leq c(T) \left( \|\mathbf{f}\|_{L_p(Q_T^1)} + \|f\|_{W_p^{1,0}(Q_T^1)} + \|\mathbf{F}\|_{W_p^{0,1}(Q_T^1)} \right. \\ &+ \|\mathbf{d}_\tau\|_{W_2^{1-1/p,1/2-1/(2p)}(G_T)} + \|\mathbf{d} \cdot \mathbf{N}\|_{W_2^{1-1/p,0}(G_T)} \\ &\left. + \|g\|_{W_p^{2-1/p,1-1/2p}(G_T)} + \|\mathbf{v}_0\|_{W_p^{2-2/p}(\mathcal{F}_1)} + \|\rho_0\|_{W_p^{3-2/p}(\mathcal{G})} \right). \end{aligned} \quad (2.5)$$

The constant  $c(T)$  in (2.5) is an increasing functions of  $T$ .

The condition  $\nabla \cdot \mathbf{v}_0(x) = f(x, 0)$ ,  $x \in \mathcal{F}_1$  can be understood in a weak sense as  $\int_{\mathcal{F}_1} (\mathbf{v}_0(y) - \mathbf{F}(y, 0)) \cdot \nabla \eta(y) dy = 0$  for arbitrary smooth  $\eta$  such that  $\eta|_{\mathcal{G}} = 0$ .

The theorem is proved in [10].

**Theorem 3.** Assume that  $\mathbf{k} = \operatorname{rot} \mathbf{K}(y, t)$ ,  $\mathbf{a} = [\mathbf{A}]$ ,  $\mathbf{K}, \mathbf{A} \in W_p^{2,1}(Q_T^i)$ ,  $i = 1, 2, 3$ ,

$$\begin{aligned} \mathbf{K} = \mathbf{A} = 0, \quad y \in S, \quad \mathbf{A} = 0, \quad y \in \mathcal{F}_3, \quad [\mathbf{K}_\tau] - [\mathbf{A}_\tau] = 0, \quad y \in S_3, \\ [\mathbf{K}_\tau] = \mathbf{a}, \quad \mathbf{A}^{(1)} \cdot \mathbf{N}(y) = \mathbf{A}^{(2)} \cdot \mathbf{N} = 0, \quad y \in \mathcal{G}. \end{aligned} \quad (2.6)$$

Then the problem (2.3) has a unique solution  $\mathbf{h} \in W_p^{2,1}(\mathcal{F}_i)$ ,  $i = 1, 2, 3$ , orthogonal to the space  $\tilde{U}_n(\Omega)$ :

$$\int_{\Omega} \mu \mathbf{h} \cdot \tilde{\mathbf{u}}_q dx = 0, \quad q = 1, \dots, b_1(\Omega). \quad (2.7)$$

The solution satisfies the inequality

$$\sum_{i=1}^3 \|\mathbf{h}\|_{W_p^{2,1}(\mathcal{F}_i)} \leq c \sum_{i=1}^3 (\|\mathbf{K}\|_{W_p^{2,1}(Q_T^i)} + \|\mathbf{A}\|_{W_p^{2,1}(Q_T^i)}). \quad (2.8)$$

**Proof.** We restrict ourselves with the formula for the solution of the problem (2.3):

$$\mathbf{h} = \mathbf{A} + \nabla\phi + \mathbf{X} + \sum_{j=1}^{b_1(\Omega)} c_j \tilde{\mathbf{u}}_j.$$

It differs from a similar formula in [3], Theorem 3, only by the last term. The functions  $\phi$ ,  $\mathbf{X}$  are the same as in [3]; they solve the problems

$$\begin{cases} \nabla^2 \phi(x) = -\nabla \cdot \mathbf{A}(x), & x \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3, & \frac{\partial \phi}{\partial N} = 0, & x \in S, \\ [\phi] = 0, & [\mu \frac{\partial \phi}{\partial n}] = 0, & x \in \mathcal{G}, & [\phi] = 0, & [\mu \frac{\partial \phi}{\partial n}] = 0, & x \in S_3, \end{cases} \quad (2.9)$$

$$\begin{cases} \operatorname{rot} \mathbf{X} = \operatorname{rot}(\mathbf{K}(x, t) - \mathbf{A}(x, t)), & \nabla \cdot \mathbf{X} = 0, & x \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3, & \mathbf{X} \cdot \mathbf{N} = 0, & x \in S, \\ [\mu \mathbf{X} \cdot \mathbf{N}] = 0, & [\mathbf{X}_\tau] = 0, & x \in \mathcal{G}, & [\mu \mathbf{X} \cdot \mathbf{n}] = 0, & [\mathbf{X}_\tau] = 0, & x \in S_3, \end{cases}$$

$$\mathbf{X}(x) = \mathbf{X}_1(x) + \nabla U(x),$$

$$\mathbf{X}_1(x) = \frac{1}{4\pi} \operatorname{rot} \int_{\Omega} \frac{\operatorname{rot}(\mathbf{K}(y) - \mathbf{A}(y))}{|x - y|} dy,$$

$$\begin{cases} \nabla^2 U(x) = 0, & x \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3, & [U(x)] = 0, & [\mu \frac{\partial U}{\partial N}] = -[\mu] \mathbf{X}_1 \cdot \mathbf{N}, & x \in \mathcal{G}, \\ [U(x)] = 0, & [\mu \frac{\partial U}{\partial n}] = -[\mu] \mathbf{X}_1 \cdot \mathbf{n}, & x \in S_3, & \frac{\partial U}{\partial n} = -\mathbf{X}_1 \cdot \mathbf{n}, & x \in S, \end{cases}$$

and they are estimated as in [3]. The orthogonality condition for  $\mathbf{h}$  yields the formulas for  $c_j(t)$ : if  $\int_{\Omega} \mu \tilde{\mathbf{u}}_j \cdot \tilde{\mathbf{u}}_q dy = \delta_{jq}$ ,  $j, q = 1, \dots, b_1(\mathcal{F}_2)$ , then

$$0 = \int_{\Omega} \mu \mathbf{h} \cdot \tilde{\mathbf{u}}_q dy = \int_{\Omega} \mu (\mathbf{A} + \nabla\phi + \mathbf{X}) \cdot \tilde{\mathbf{u}}_q dy + c_q(t),$$

This formula and the estimates of  $\phi$ ,  $\mathbf{X}$  (see [3]) imply (2.8).

The uniqueness of the solution follows from the fact that in the case  $\mathbf{K} = 0$ ,  $\mathbf{A} = 0$  the solution belongs to  $\tilde{U}_n(\Omega)$ . The theorem is proved.

**Theorem 4.** Assume that the data of the problem (2.2) possess the following properties:  $\mathbf{G} \in L_p(Q_T^1)$ ,  $\mathbf{H}_0 \in W_p^{2-2/p}(\mathcal{F}_j)$ ,  $j = 1, 2, 3$ ,  $\boldsymbol{\ell} \in W_p^{2,1}(Q_T^2)$ ,  $\boldsymbol{\ell}|_{x \in S} = 0$ ,  $\mathbf{a} \in W_p^{2-1/p, 1-1/2p}(G_T)$ , moreover,  $\mathbf{a} = [\mathbf{A}]$  with  $\mathbf{A}^{(i)} \in W_p^{2,1}(Q_T^i)$ ,  $i = 1, 2$ , satisfying (2.6), finally, let the compatibility conditions

$$\begin{aligned} \nabla \cdot \mathbf{G}(x, t) &= 0, & x \in \mathcal{F}_1 \cup \mathcal{F}_3, \\ \nabla \cdot \mathbf{H}_0(x) &= 0, & x \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3, & \operatorname{rot} \mathbf{H}_0(x) = \operatorname{rot} \boldsymbol{\ell}(x, 0), & x \in \mathcal{F}_2, \\ [\mu \mathbf{H}_0 \cdot \mathbf{n}] &= 0, & [\mathbf{H}_{0\tau}] = 0, & x \in S_3, \\ [\mu \mathbf{H}_0 \cdot \mathbf{N}] &= 0, & [\mathbf{H}_{0\tau}] = \mathbf{a}(x, 0) = [\mathbf{A}(x, 0)], & x \in \mathcal{G}, & \mathbf{H}_0 \cdot \mathbf{n} = 0, & x \in S \end{aligned} \quad (2.10)$$

hold. Then the problem (2.2) has a unique solution  $\mathbf{H} \in W_p^{2,1}(Q_T^i)$ ,  $i = 1, 2, 3$ , and

$$\begin{aligned} \sum_{i=1}^3 \|\mathbf{H}\|_{W_p^{2,1}(Q_T^i)} &\leq c \left( \sum_{i=1,3} \|\mathbf{G}\|_{L_p(Q_T^1)} + \sum_{i=1}^3 \|\mathbf{H}_0\|_{W_p^{2-2/p}(\mathcal{F}_i)} \right. \\ &\quad \left. + \|\boldsymbol{\ell}\|_{W_p^{2,1}(Q_T^2)} + \sum_{i=1}^2 \|\mathbf{A}\|_{W_p^{2,1}(Q_T^i)} + \sum_{q=1}^{b_1(\mathcal{F}_2)} \|\mathbf{M}_q\|_{L_p(0,T)} \right) \end{aligned} \quad (2.11)$$

**Proof.** We reduce (2.2) to a similar problem with  $\ell = 0$ ,  $\mathbf{a} = 0$ . We extend  $\ell$  into  $\Omega$  with the preservation of class, i.e., so that the extended function  $\ell^* \in W_p^{2,1}(Q_T)$  satisfies

$$\|\ell^*\|_{W_p^{2,1}(Q_T)} \leq c\|\ell\|_{W_p^{2,1}(Q_T^2)}, \quad (2.12)$$

and we define

$$\begin{aligned} \mathbf{a}^*(x, t) &= \mathbf{A}^{(1)}(x, t) - \mathbf{A}^{(2)*}(x, t), \quad x \in \mathcal{F}_1, \\ \mathbf{a}^*(x, t) &= 0, \quad x \in \mathcal{F}_2 \cup \mathcal{F}_3, \end{aligned}$$

where  $\mathbf{A}^{(2)*}$  is the extension of  $\mathbf{A}^{(2)}$  such that

$$\|\mathbf{A}^{(2)*}\|_{W_p^{2,1}(Q_T^1 \cup Q_T^2)} \leq c\|\mathbf{A}^{(2)}\|_{W_p^{2,1}(Q_T^2)}.$$

We set  $\mathbf{A}^{(3)} = 0$ . It is easily verified that

$$[\mathbf{a}^*] = [\mathbf{A}] = \mathbf{a}, \quad [\ell^* + \mathbf{a}^*] = \mathbf{a}, \quad x \in \mathcal{G}.$$

Now we define  $\mathbf{h}_1(x, t)$  as a solution of the problem (2.3) with  $\mathbf{k}(x, t) = \text{rot}(\ell^* + \mathbf{a}^*)$ . By (2.8),

$$\sum_{i=1}^3 \|\mathbf{h}_1\|_{W_p^{2,1}(Q_T^i)} \leq c(\|\ell\|_{W_p^{2,1}(Q_T^2)} + \sum_{i=1}^2 \|\mathbf{A}\|_{W_p^{2,1}(Q_T^i)}). \quad (2.13)$$

For  $\mathbf{h} = \mathbf{H} - \mathbf{h}_1$  we obtain the problem

$$\begin{cases} \mu \mathbf{h}_t(y, t) + \alpha^{-1} \text{rot} \text{rot} \mathbf{h}(y, t) = \mathbf{G}'(y, t), & \nabla \cdot \mathbf{h}(y, t) = 0, \quad y \in \mathcal{F}_1 \cup \mathcal{F}_3, \\ \text{rot} \mathbf{h}(y, t) = 0, & \nabla \cdot \mathbf{h}(y, t) = 0, \quad y \in \mathcal{F}_2 \\ [\mu \mathbf{h} \cdot \mathbf{N}] = 0, & [\mathbf{h}_\tau] = 0, \quad y \in \mathcal{G}, \\ [\mu \mathbf{h} \cdot \mathbf{n}] = 0, & [\mathbf{h}_\tau] = 0, \quad y \in S_3, \quad \mathbf{h} \cdot \mathbf{n} = 0, \quad y \in S, \\ \mathbf{h}(y, 0) = \mathbf{h}_0(y) = \mathbf{H}_0(y) - \mathbf{h}_1(y, 0), & y \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3, \end{cases} \quad (2.14)$$

$$\begin{cases} \int_{\mathcal{F}_2} \mu \mathbf{h}_t \cdot \mathbf{u}_q^\mathcal{F} dy = \int_{\mathcal{G}} (\mathbf{N} \times \alpha^{-1} \text{rot} \mathbf{h}^{(1)}) \cdot \mathbf{u}_q^\mathcal{F}(y) dS \\ \quad + \int_{S_3} (\mathbf{n} \times \alpha^{-1} \text{rot} \mathbf{h}^{(3)}) \cdot \mathbf{u}_q^\mathcal{F}(y) dS + \mathbf{M}'_q(t), \quad q = 1, \dots, b_1(\mathcal{F}_2), \end{cases} \quad (2.15)$$

where

$$\begin{aligned} \mathbf{G}' &= \mathbf{G} - \mu \mathbf{h}_{1t} - \alpha^{-1} \text{rot} \text{rot} \mathbf{h}_1, \\ \mathbf{M}'_q &= \mathbf{M}_q + \int_{\mathcal{G}} (\mathbf{N} \times \text{rot} \mathbf{h}_1^{(1)}) \cdot \mathbf{u}_q^\mathcal{F}(y) dS \\ &\quad + \int_{S_3} (\mathbf{n} \times \text{rot} \mathbf{h}_1^{(3)}) \cdot \mathbf{u}_q^\mathcal{F}(y) dS - \int_{\mathcal{F}_2} \mu \mathbf{h}_{1t} \cdot \mathbf{u}_q^\mathcal{F} dy, \quad q = 1, \dots, b_1(\mathcal{F}_2). \end{aligned}$$

The equations  $\text{rot} \mathbf{h}(y, t) = 0$ ,  $\nabla \cdot \mathbf{h} = 0$ ,  $y \in \mathcal{F}_2$  imply

$$\mathbf{h}^{(2)} = \nabla \varphi(y, t) + \sum_{j=1}^{b_1(\mathcal{F}_2)} k_j(t) \mathbf{u}_j^\mathcal{F}(y), \quad y \in \mathcal{F}_2, \quad (2.16)$$

where  $\varphi$  is a solution to the problem

$$\begin{cases} \nabla^2 \varphi(y, t) = 0, & y \in \mathcal{F}_2, & \frac{\partial \varphi}{\partial n} = 0, & y \in S, \\ \mu_2 \frac{\partial \varphi}{\partial N} = \mu_1 \mathbf{h}^{(1)} \cdot \mathbf{N}, & y \in \mathcal{G}, & \mu_2 \frac{\partial \varphi}{\partial n} = \mu_3 \mathbf{h}^{(3)} \cdot \mathbf{n}, & y \in S_3. \end{cases} \quad (2.17)$$

It follows that  $\mathbf{h}^{(1)}$ ,  $\mathbf{h}^{(3)}$  and  $\varphi$  satisfy

$$\begin{cases} \mu \mathbf{h}_t(y, t) + \alpha^{-1} \text{rot} \text{rot} \mathbf{h}(y, t) = \mathbf{G}'(y, t), & \nabla \cdot \mathbf{h}(y, t) = 0, & y \in \mathcal{F}_1 \cup \mathcal{F}_3, \\ \nabla^2 \varphi(y, t) = 0, & y \in \mathcal{F}_2, & \frac{\partial \varphi}{\partial n} = 0, & y \in S, \\ \mu_2 \frac{\partial \varphi}{\partial N} = \mu_1 \mathbf{h}^{(1)} \cdot \mathbf{N}, & y \in \mathcal{G}, & \mu_2 \frac{\partial \varphi}{\partial n} = \mu_3 \mathbf{h}^{(3)} \cdot \mathbf{n}, & y \in S_3, \\ \mathbf{h}_\tau^{(1)} - \nabla_\tau \varphi = \sum_{j=1}^{b_1(\mathcal{F}_2)} k_j(t) \mathbf{u}_j^\mathcal{F}(y), & y \in \mathcal{G}, \\ \mathbf{h}_\tau^{(3)} - \nabla_\tau \varphi = \sum_{j=1}^{b_1(\mathcal{F}_2)} k_j(t) \mathbf{u}_j^\mathcal{F}(y), & y \in S_3, \\ \mathbf{h}(y, 0) = \mathbf{h}_0(y) = \mathbf{H}_0(y) - \mathbf{h}_1(y, 0), & y \in \mathcal{F}_1 \cup \mathcal{F}_3. \end{cases} \quad (2.18)$$

This problem is studied in [11] under the assumptions  $k_j = 0$ ,  $S_3 = \emptyset$ , but the result remains valid in the case considered here, in particular, there holds the a-priori estimate

$$\begin{aligned} & \sum_{i=1,3} \|\mathbf{h}^{(i)}\|_{W_p^{2,1}(Q_T^i)} + \|\nabla \varphi\|_{W_p^{2,1}(Q_T^2)} \\ & \leq c \left( \sum_{i=1,3} (\|\mathbf{G}'\|_{L_p(Q_T^i)} + \|\mathbf{h}_0\|_{W_p^{2-1/p}(\mathcal{F}_i)}) + \|\mathbf{k}\|_{W_p^{1-1/2p}(0,T)} \right). \end{aligned} \quad (2.19)$$

The functions  $k_j(t)$  can be expressed in terms of  $\mathbf{h}$  with the help of equations (2.15), (2.16): if  $\int_{\mathcal{F}_2} \mu \mathbf{u}_m^\mathcal{F} \cdot \mathbf{u}_q^\mathcal{F} dy = \delta_{mq}$ ,  $q, m = 1, \dots, b_1(\mathcal{F}_2)$ , then (2.15) imply

$$\begin{aligned} k_j(t) &= k_j(0) + \int_0^t \mathbf{M}'_j(\tau) d\tau + \int_0^t d\tau \left( \int_{\mathcal{G}} (\mathbf{N} \times \text{rot} \mathbf{h}^{(1)}) \cdot \mathbf{u}_j^\mathcal{F} dS \right. \\ & \quad \left. + \int_{S_3} (\mathbf{n} \times \text{rot} \mathbf{h}^{(3)}) \cdot \mathbf{u}_j^\mathcal{F} dS \right), \quad j = 1, \dots, b_1(\mathcal{F}_2), \end{aligned} \quad (2.20)$$

and

$$\begin{aligned} \|\mathbf{k}\|_{W_p^1(0,T)} &\leq c(T) (|\mathbf{k}(0)| + \sum_q \|\mathbf{M}'_q\|_{L_p(0,T)} \\ & \quad + \|\text{rot} \mathbf{h}^{(1)}\|_{L_p(G_T)} + \|\text{rot} \mathbf{h}^{(3)}\|_{L_p(S_3 \times (0,T))} + \|\mathbf{k}\|_{W_p^{1-1/2p}(0,T)}). \end{aligned} \quad (2.21)$$

The norms of  $\mathbf{h}$  in (2.21) can be estimated by (2.19) and the interpolation inequalities, which yields

$$\begin{aligned} & \sum_{i=1,3} \|\mathbf{h}^{(i)}\|_{W_p^{2,1}(Q_T^i)} + \|\nabla \varphi\|_{W_p^{2,1}(Q_T^2)} + \|\mathbf{k}\|_{W_p^1(0,T)} \\ & \leq c(T) \left( \sum_{i=1,3} \|\mathbf{G}'\|_{L_p(Q_T^i)} + \sum_{i=1}^3 \|\mathbf{h}_0\|_{W_p^{2-1/p}(\mathcal{F}_i)} + \sum_q \|\mathbf{M}'_q\|_{L_p(0,T)} + \|\mathbf{h}^{(i)}\|_{L_p(Q_T)} \right. \\ & \quad \left. + \|\mathbf{k}\|_{L_p(0,T)} \right). \end{aligned} \quad (2.22)$$

This inequality holds for arbitrary  $T' < T$ , hence, by the Gronwall lemma,

$$\begin{aligned} & \sum_{i=1,3} \|\mathbf{h}^{(i)}\|_{W_p^{2,1}(Q_T^i)} + \|\nabla \varphi\|_{W_p^{2,1}(Q_T^2)} + \|\mathbf{k}\|_{W_p^1(0,T)} \\ & \leq c(T) \left( \sum_{i=1,3} \|\mathbf{G}'\|_{L_p(Q_T^i)} + \sum_{i=1}^3 \|\mathbf{h}_0\|_{W_p^{2-1/p}(\mathcal{F}_i)} + \sum_q \|\mathbf{M}'_q\|_{L_p(0,T)} \right) \end{aligned} \quad (2.23)$$

Finally, the estimate of  $\mathbf{h}^{(2)}$  is obtained from (2.16). Taking (2.13) into account, we arrive at (2.11).

Our arguments show that the term  $\sum_{j=1}^{b_1(\mathcal{F}_2)} k_j(t) \mathbf{u}_j(y)$  in (2.18) generates a compact operator, which makes it possible to deduce the solvability of the problem (2.18), (2.15) from the solvability of the same problem with  $k_j = 0$  (established in [11]) by standard methods. Theorem 4 is proved.

### 3 Nonlinear problem

In this section we outline main ideas of the proof of Theorem 1. We consider the problem (1.6), (1.15), (1.16) that can be written in the form

$$\left\{ \begin{aligned} & \mathbf{u}_t(y, t) - \nu \nabla^2 \mathbf{u} + \nabla q = \mathbf{l}_1(\mathbf{u}, q, \mathbf{h}, \rho) + \mathbf{f}(y, t), \\ & \nabla \cdot \mathbf{u} = l_2(\mathbf{u}, \rho), \quad y \in \mathcal{F}_1, \quad t > 0, \\ & \Pi_{\mathcal{G}} S(\mathbf{u}) \mathbf{N} = \mathbf{l}_3(\mathbf{u}, \rho), \\ & -q + \nu \mathbf{N} \cdot S(\mathbf{u}) \mathbf{N}(y) + \sigma \mathfrak{B}(\rho) = l_4(\mathbf{u}, \mathbf{h}, \rho) + l_5(\rho) + \sigma \mathcal{H}(y), \\ & \rho_t + \mathbf{V}(x) \cdot \nabla_{\tau} \rho - \mathbf{u} \cdot \mathbf{N}(y) = l_6(\mathbf{u}, \rho), \quad y \in \mathcal{G}, \\ & \mathbf{u}(y, 0) = \mathbf{u}_0(y), \quad y \in \mathcal{F}_1, \quad \rho(y, 0) = \rho_0(y), \quad y \in \mathcal{G}, \end{aligned} \right. \quad (3.1)$$

$$\left\{ \begin{aligned} & \mu \mathbf{h}_t + \alpha^{-1} \text{rot} \text{rot} \mathbf{h} = \mathbf{l}_7(\mathbf{h}, \mathbf{u}, \rho), \\ & \nabla \cdot \mathbf{h} = 0, \quad y \in \mathcal{F}_1, \\ & \mu \mathbf{h}_t + \alpha^{-1} \text{rot} \text{rot} \mathbf{h} = \alpha^{-1} \text{rot} \mathbf{j}(y, t), \\ & \nabla \cdot \mathbf{h} = 0, \quad y \in \mathcal{F}_3, \\ & \text{rot} \mathbf{h} = \text{rot} \mathbf{l}_8(\mathbf{h}, \rho), \quad \nabla \cdot \mathbf{h} = 0, \quad y \in \mathcal{F}_2, \\ & [\mu \mathbf{h} \cdot \mathbf{N}] = 0, \quad [\mathbf{h}_{\tau}] = \mathbf{l}_9(\mathbf{h}, \rho), \quad y \in \mathcal{G}, \\ & [\mu \mathbf{h} \cdot \mathbf{n}] = 0, \quad [\mathbf{h}_{\tau}] = 0, \quad y \in S_3, \\ & \mathbf{h} \cdot \mathbf{n} = 0, \quad y \in S, \\ & \mathbf{h}(y, 0) = \mathbf{h}_0(y), \quad y \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3, \end{aligned} \right. \quad (3.2)$$

$$\left\{ \begin{aligned} & \int_{\mathcal{F}_2} \mu \mathbf{h}_t \cdot \mathbf{u}_q^{\mathcal{F}} dy - \int_{\mathcal{G}} \mathbf{N} \times \alpha^{-1} \text{rot} \mathbf{h}^{(1)} \cdot \mathbf{u}_q^{\mathcal{F}} dS - \int_{S_3} \mathbf{n} \times \alpha^{-1} \text{rot} \mathbf{h}^{(3)} \cdot \mathbf{u}_q^{\mathcal{F}} dS \\ & = l_{10,q}(\mathbf{u}, \rho, \mathbf{h}) - \int_{S_3} \alpha^{-1} \mathbf{j} \cdot \mathbf{u}_q^{\mathcal{F}} dS, \quad q = 1, \dots, b_1(\mathcal{F}_2), \end{aligned} \right. \quad (3.3)$$

where

$$\left\{ \begin{aligned}
& l_1(\mathbf{u}, q, \rho) = \nu(\tilde{\nabla}^2 - \nabla^2)\mathbf{u} + (\nabla - \tilde{\nabla})q + \rho_t^*(\mathcal{L}^{-1}(y, \rho^*)\mathbf{N}^*(y) \cdot \nabla)\mathbf{u} \\
& \quad - (\mathcal{L}^{-1}\mathbf{u} \cdot \nabla)\mathbf{u} + \tilde{\nabla} \cdot T_M(\frac{\mathcal{L}}{L}\mathbf{h}) + \int_0^1 \rho^*(y, t)(\mathbf{N}^*(y) \cdot \nabla_x)\mathbf{f}(x, t)|_{x=e_{s\rho}} ds, \\
& l_2(\mathbf{u}, \rho) = (I - \hat{\mathcal{L}}^T)\nabla \cdot \mathbf{u} = \nabla \cdot \mathbf{L}(\mathbf{u}, \rho), \quad \mathbf{L} = (I - \hat{\mathcal{L}})\mathbf{u}, \quad y \in \mathcal{F}_1, \\
& l_3(\mathbf{u}, \rho) = \Pi_{\mathcal{G}}(\Pi_{\mathcal{G}}S(\mathbf{u})\mathbf{N})(y) - \Pi\tilde{S}(\mathbf{u})\mathbf{n}(e_{\rho}(y)), \\
& l_4(\mathbf{u}, \mathbf{h}, \rho) = \nu(\mathbf{N} \cdot S(\mathbf{u})\mathbf{N} - \mathbf{n} \cdot \tilde{S}(\mathbf{u})\mathbf{n}) - [T_M(\frac{\mathcal{L}}{L}\mathbf{h})]\mathbf{n}, \\
& l_5(\rho) = - \int_0^1 (1-s) \frac{d^2}{ds^2} \mathcal{L}^{-T}(y, s\rho) \nabla \cdot \frac{\mathcal{L}^T(y, s\rho)\mathbf{N}}{|\mathcal{L}^T(y, s\rho)\mathbf{N}|} ds, \\
& l_6(\mathbf{u}, \mathbf{h}, \rho) = (\frac{\hat{\mathcal{L}}^T\mathbf{N}}{|\hat{\mathcal{L}}^T\mathbf{N}|} + \nabla_{\tau}\rho - \mathbf{N}) \cdot \mathbf{u} + (\mathbf{V} - \mathbf{u}) \cdot \nabla_{\tau}\rho, \quad y \in \mathcal{G}, \\
& l_7(\mathbf{h}, \rho) = \alpha^{-1}\text{rot}(\text{rot}\mathbf{h} - \mathcal{P}\text{rot}\mathcal{P}\mathbf{h}) + \frac{1}{L}\hat{\mathcal{L}}_t\mathcal{L}\mathbf{h} \\
& \quad + \rho_t^*\hat{\mathcal{L}}(\mathcal{L}^{-1}\mathbf{N} \cdot \nabla)\frac{1}{L}\mathcal{L}\mathbf{h} + \mu_1\text{rot}(\mathcal{L}^{-1}\mathbf{u} \times \mathbf{h}), \quad y \in \mathcal{F}_1, \\
& l_8(\mathbf{h}, \rho) = (I - \mathcal{P})\mathbf{h}, \quad y \in \mathcal{F}_2, \\
& l_9(\mathbf{h}, \rho) = (\frac{\hat{\mathcal{L}}\hat{\mathcal{L}}^T\mathbf{N}}{|\hat{\mathcal{L}}^T\mathbf{N}|^2} - \mathbf{N})[\mathbf{h} \cdot \mathbf{N}] = [\mathbf{A}(\mathbf{h}, \rho)], \quad y \in \mathcal{G}, \\
& \mathbf{A}^{(i)} = (\frac{\hat{\mathcal{L}}(y, \rho^*)\hat{\mathcal{L}}^T\mathbf{N}^*(y)}{|\hat{\mathcal{L}}^T\mathbf{N}^*|^2} - \frac{\mathbf{N}^*(y)}{|\mathbf{N}^*|^2})(\mathbf{N}^* \cdot \mathbf{h}^{(i)}), \quad i = 1, 2, \quad \mathbf{A}^{(3)} = 0, \\
& l_{10,q}(\mathbf{u}, \rho, \mathbf{h}) = \int_{\mathcal{F}_2} \mu\Phi \cdot \mathbf{u}_q^{\mathcal{F}} dy + \int_{\mathcal{G}} \mathbf{N} \times \alpha^{-1}(\mathcal{P}\text{rot}\mathcal{P}\mathbf{h}^{(1)} - \text{rot}\mathbf{h}^{(1)}) \cdot \mathbf{u}_q^{\mathcal{F}} dS \\
& \quad - \int_{\mathcal{G}} (\mathbf{N} \times \mu(\mathcal{L}^{-1}\mathbf{u} \times \mathbf{h}^{(1)}) - \Psi) \cdot \mathbf{u}_q^{\mathcal{F}} dS, \quad q = 1, \dots, b_1(\mathcal{F}_2).
\end{aligned} \right. \tag{3.4}$$

We have used the formula for the variation of the mean curvature under the normal perturbation of the surface:

$$H(e_{\rho}) - \mathcal{H}(y) = -\mathfrak{B}\rho - \int_0^1 (1-s) \frac{d^2}{ds^2} \mathcal{L}^{-T}(y, s\rho) \nabla \cdot \frac{\mathcal{L}^T(y, s\rho)\mathbf{N}}{|\mathcal{L}^T(y, s\rho)\mathbf{N}|} ds.$$

and we have set

$$\Pi_{\mathcal{G}}\mathbf{a} = \mathbf{a} - \mathbf{N}(\mathbf{a} \cdot \mathbf{N}), \quad \Pi\mathbf{a} = \mathbf{a} - \mathbf{n}(\mathbf{a} \cdot \mathbf{n}).$$

The vector field  $\mathbf{V} \in W_p^{2-1/p}(\mathcal{G})$  is introduced to optimize the estimate of the expression  $l_6$  in (3.4).

The solvability of the problem (3.1)-(3.4) is proved by successive approximations, according to a standard scheme:

$$\left\{ \begin{aligned}
& \mathbf{u}_{m+1,t}(y, t) - \nu\nabla^2\mathbf{u}_{m+1} + \nabla q_{m+1} = l_1(\mathbf{u}_m, q_m, \mathbf{h}_m, \rho_m) + \mathbf{f}(y, t), \\
& \nabla \cdot \mathbf{u}_{m+1} = l_2(\mathbf{u}_m, \rho_m), \quad y \in \mathcal{F}_1, \quad t > 0, \\
& \Pi_{\mathcal{G}}S(\mathbf{u}_{m+1})\mathbf{N} = l_3(\mathbf{u}_m, \rho_m), \\
& -q_{m+1} + \nu\mathbf{N} \cdot S(\mathbf{u}_{m+1})\mathbf{N}(y) + \sigma\mathfrak{B}(\rho_{m+1}) = l_4(\mathbf{u}_m, \mathbf{h}_m, \rho_m) + l_5(\rho_m) + \sigma\mathcal{H}(y), \\
& \rho_{m+1,t} + \mathbf{V}(x) \cdot \nabla_{\tau}\rho_{m+1} - \mathbf{u}_{m+1} \cdot \mathbf{N}(y) = l_6(\mathbf{u}_m, \rho_m), \quad y \in \mathcal{G}, \\
& \mathbf{u}_{m+1}(y, 0) = \mathbf{u}_0(y), \quad y \in \mathcal{F}_1, \quad \rho_{m+1}(y, 0) = \rho_0(y), \quad y \in \mathcal{G},
\end{aligned} \right. \tag{3.5}$$



$$\left\{ \begin{array}{l}
\mu \mathbf{h}_{m+1,t} + \alpha^{-1} \text{rot} \text{rot} \mathbf{h}_{m+1} = \mathbf{l}_7(\mathbf{h}_m, \mathbf{u}_m, \rho_m), \\
\nabla \cdot \mathbf{h}_{m+1} = 0, \quad y \in \mathcal{F}_1, \\
\mu \mathbf{h}_{m+1,t} + \alpha^{-1} \text{rot} \text{rot} \mathbf{h}_{m+1} = \alpha^{-1} \text{rot} \mathbf{j}(y, t), \\
\nabla \cdot \mathbf{h}_{m+1} = 0, \quad y \in \mathcal{F}_3, \\
\text{rot} \mathbf{h}_{m+1} = \text{rot} \mathbf{l}_8(\mathbf{h}_m, \rho_m), \quad \nabla \cdot \mathbf{h}_{m+1} = 0, \quad y \in \mathcal{F}_2, \\
[\mu \mathbf{h}_{m+1} \cdot \mathbf{N}] = 0, \quad [\mathbf{h}_{m+1,\tau}] = \mathbf{l}_9(\mathbf{h}_m, \rho_m), \quad y \in \mathcal{G}, \\
[\mu \mathbf{h}_{m+1} \cdot \mathbf{n}] = 0, \quad [\mathbf{h}_{m+1,\tau}] = 0, \quad y \in S_3, \\
\mathbf{h}_{m+1} \cdot \mathbf{n} = 0, \quad y \in S, \\
\int_{\mathcal{F}_2} \mu \mathbf{h}_{m+1,t} \cdot \mathbf{u}_q^{\mathcal{F}} dy - \int_{\mathcal{G}} \mathbf{N} \times \alpha^{-1} \text{rot} \mathbf{h}_{m+1} \cdot \mathbf{u}_q^{\mathcal{F}} dS - \int_{S_3} \mathbf{n} \times \alpha^{-1} \text{rot} \mathbf{h}_{m+1} \cdot \mathbf{u}_q^{\mathcal{F}} dS \\
= l_{10,q}(\mathbf{u}_m, \rho_m, \mathbf{h}_m) - \int_{S_3} \mathbf{j} \cdot \mathbf{u}_q^{\mathcal{F}} dS, \quad q = 1, \dots, b_1(\mathcal{F}_2), \\
\mathbf{h}_{m+1}(y, 0) = \mathbf{h}_0(y), \quad y \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3.
\end{array} \right. \quad (3.6)$$

The first approximation,  $(\mathbf{u}_1, q_1, \rho_1, \mathbf{h}_1)$ , is defined as follows:  $q_1 = 0$ ,  $\mathbf{u}_1, \rho_1, \mathbf{h}_1$  satisfy  $\mathbf{u}_1(y, 0) = \mathbf{u}_0(y)$ ,  $y \in \mathcal{F}_1$ ,  $\rho_1(y, 0) = \rho_0(y)$ ,  $y \in \mathcal{G}$ ,  $\mathbf{h}_1(y, 0) = \mathbf{h}_0(y)$ ,  $y \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3$  and the inequalities

$$\begin{aligned}
\|\mathbf{u}_1\|_{W_p^{2,1}(Q_T^1)} &\leq c \|\mathbf{u}_1\|_{W_p^{2,1}(Q_\infty^1)} \leq c \|\mathbf{u}_0\|_{W_p^{2-2/p}(\mathcal{F}_1)}, \\
\|\rho_1\|_{W_p^{3-1/p,0}(G_\infty)} + \|\rho_{1,t}\|_{W_p^{2-1/p,1-1/2p}(G_\infty)} &\leq c \|\rho_0\|_{W_p^{3-2/p}(\mathcal{G})}, \\
\sum_{i=1}^2 \|\mathbf{h}_1^{(i)}\|_{W_r^{2,1}(Q_T^i)} &\leq c \|\mathbf{h}_0^{(1)}\|_{W_r^{2-2/r}(\mathcal{F}_1)};
\end{aligned} \quad (3.7)$$

in addition,  $\mathbf{h}_1$  is divergence free and satisfies (2.16), (2.17) (for the construction of such  $\mathbf{h}_1$ , see Theorem 4 in [2]).

The proof of the solvability of (3.4) is based on Theorems 2, 4 (with the exponent  $r$  instead of  $p$ ) and on the estimates of nonlinear terms (3.2) obtained in [3], Sec. 5.

**Theorem 5.** Assume that  $\mathbf{u}_m \in W_p^{2,1}(Q_T^1)$ ,  $\nabla q_m \in L_p(Q_T^1)$ ,  $\rho_m \in W_p^{3-1/p,0}(G_T)$ ,  $\rho_{m,t} \in W_p^{2-1/p,1-1/2p}(G_T)$ ,  $\mathbf{h}_m \in W_r^{2,1}(Q_T^i)$ ,  $i = 1, 2, 3$ ,  $\mathbf{V} \in W_p^{2-1/p}(\mathcal{G})$ , and the conditions (1.19), (1.22), as well as

$$\begin{aligned}
\|\rho_m(\cdot, t)\|_{W_p^{1-1/p}(\mathcal{G})} &\leq \delta \ll 1, \\
\|\mathbf{V} - \mathbf{u}_0\|_{W_p^{1-1/p}(\mathcal{G})} &\leq \delta, \\
T^{1/2-1/p} \|\mathbf{V}\|_{W_p^{2-1/p}(\mathcal{G})} &\leq \delta
\end{aligned} \quad (3.8)$$

are satisfied. Then

$$Z_m(T) \leq \vartheta(\delta, T)(Y_m(T) + Y_m^2(T)), \quad (3.9)$$

where

$$\begin{aligned}
Z_m(T) = & \|l_1(\mathbf{u}_m, q_m, \mathbf{h}_m, \rho)\|_{L_p(Q_T^1)} + \|l_2(\mathbf{u}_m, \rho_m)\|_{W_p^{1,0}(Q_T^1)} + \|\mathbf{L}(\mathbf{u}_m, \rho_m)\|_{W_p^{0,l}(Q_T^1)} \\
& + \|l_3(\mathbf{u}_m, \rho_m)\|_{W_p^{1-1/p, 1/2-1/2p}(G_T)} + \|l_4(\mathbf{u}_m, \mathbf{h}_m, \rho_m)\|_{W_p^{1-1/p, 0}(G_T)} \\
& + \|l_5(\rho_m)\|_{W_p^{1-1/p, 0}(G_T)} + \|l_6(\mathbf{u}_m, \rho_m)\|_{W_p^{2-1/p, 1-1/2p}(G_T)} + \sum_{i=1}^3 \|l_7(\mathbf{h}, \rho)\|_{L_r(Q_T^1)} \\
& + \|l_8(\mathbf{h}_m, \rho_m)\|_{W_r^{2,0}(Q_T^2)} + \sum_{i=1}^2 \|\mathbf{A}^{(i)}\|_{W_r^{2,1}(Q_T^i)},
\end{aligned}$$

and  $\vartheta(\delta, T)$  is a constant that can be made arbitrarily small by the choice of small  $\delta$  and  $T$ .

Inequality (3.9) is established in [3], Sec. 5, except for the estimate of  $\rho^* \mathbf{N}^* \cdot \nabla \mathbf{f}|_{x=e_{s\rho}}$  that is obtained as follows:

$$\begin{aligned}
\|\rho^* \mathbf{N}^* \cdot \nabla \mathbf{f}|_{x=e_{s\rho}}\|_{L_p(Q_T)} & \leq cT^{1/p} \sup_{Q_T^1} |\rho^*(x, t)| \sup_{t < T} \|\nabla \mathbf{f}(\cdot, t)\|_{L_p(\mathbb{R}^3)} \\
& \leq cT^{1/p} \sup_{t < T} \|\rho\|_{W_p^{1-1/p}(\mathcal{G})}.
\end{aligned} \tag{3.10}$$

The nonlinear expressions in  $l_{10,q}$  are also estimated in [3].

If  $(\mathbf{u}_m, q_m, \rho_m, \mathbf{h}_m)$  are found, then, by Theorems 2, 4,5 the problem (3.5), (3.6) is uniquely solvable and

$$Y_{m+1}(T) \leq c\vartheta(\delta, T)(Y_m(T) + Y_m^2(T)) + c_1 N, \tag{3.11}$$

where

$$N = \|\mathbf{u}_0\|_{W_p^{2-2/p}(\mathcal{F}_1)} + \|\rho_0\|_{W_p^{3-2/p}(\mathcal{G})} + \|\mathcal{H}\|_{W_p^{1-1/p}(\mathcal{G})} + \sum_{i=1}^3 \|\mathbf{h}_0\|_{W_r^{2-2/r}(\mathcal{F}_i)}, \tag{3.12}$$

$$\begin{aligned}
Y_m(T) = & \|\mathbf{u}_m\|_{W_p^{2,1}(Q_T^1)} + \sup_{t < T} \|\mathbf{u}_m\|_{W_p^{2-1/p}(\mathcal{F}_1)} + \|\nabla q_m\|_{L_p(Q_T^1)} \\
& + \|q_m\|_{W_p^{1-1/p}(G_T)} + \|\rho_m\|_{W_p^{3-1/p, 0}(G_T)} + \|\rho_{m,t}\|_{W_p^{2-2/p, 1-1/2p}(G_T)} \\
& + \sum_{i=1}^2 (\|\mathbf{h}_m\|_{W_r^{2,1}(Q_T^i)} + \sup_{t < T} \|\mathbf{h}_m\|_{W_r^{2-2/r}(\mathcal{F}_i)}) \equiv Y(\mathbf{u}_m, q_m, \rho_m, \mathbf{h}_m).
\end{aligned} \tag{3.13}$$

It follows that

$$Y_{m+1}(T) \leq 2c_1 N, \tag{3.14}$$

if  $Y_m(T) \leq 2c_1 N$  and  $\vartheta$  is so small that

$$\vartheta(\delta, T)(2c_1 N + 4(c_1 N)^2) \leq c_1 N$$

i.e.,

$$\vartheta(\delta, T) \leq \frac{1}{2(1 + 2c_1 N)}; \tag{3.15}$$

in addition, for small  $\epsilon$  and  $T$ ,

$$\begin{aligned}
\|\rho_{m+1}(\cdot, t)\|_{W_p^{2-2/p}(\mathcal{G})} & \leq \|\rho_0\|_{W_p^{2-1/p}(\mathcal{G})} + \int_0^t \|\rho_{m+1,\tau}(\cdot, \tau)\|_{W_p^{2-1/p}(\mathcal{G})} d\tau \\
& \leq \epsilon + 2c_1 T^{1-1/p} N \leq \delta.
\end{aligned} \tag{3.16}$$

In the case  $m = 0$  this inequality follows from (3.7), (1.11). Hence if  $c_1$  is chosen in such a way that (3.14) holds for  $m = 0$ , then (3.14) and (3.16) are satisfied for all  $m$ .

In order to prove the convergence of the sequence  $(\mathbf{u}_m, q_m, \rho_m, \mathbf{h}_m)$ , we estimate the differences

$$\mathbf{w}_{m+1} = \mathbf{u}_{m+1} - \mathbf{u}_m, \quad s_{m+1} = q_{m+1} - q_m, \quad r_{m+1} = \rho_{m+1} - \rho_m, \quad \mathbf{k}_{m+1} = \mathbf{h}_{m+1} - \mathbf{h}_m.$$

They satisfy the relations

$$\left\{ \begin{array}{l} \mathbf{w}_{m+1,t}(y, t) - \nu \nabla^2 \mathbf{w}_{m+1} + \nabla s_{m+1} = \mathbf{l}_1(\mathbf{u}_m, q_m, \mathbf{h}_m, \rho_m) - \mathbf{l}_1(\mathbf{u}_{m-1}, q_{m-1}, \mathbf{h}_{m-1}, \rho_{m-1}), \\ \nabla \cdot \mathbf{w}_{m+1} = l_2(\mathbf{u}_m, \rho_m) - l_2(\mathbf{u}_{m-1}, \rho_{m-1}), \quad y \in \mathcal{F}_1, \quad t > 0, \\ \Pi_{\mathcal{G}} S(\mathbf{w}_{m+1}) \mathbf{N} = \mathbf{l}_3(\mathbf{u}_m, \rho_m) - \mathbf{l}_3(\mathbf{u}_{m-1}, \rho_{m-1}), \\ -s_{m+1} + \nu \mathbf{N} \cdot S(\mathbf{w}_{m+1}) \mathbf{N}(y) + \sigma \mathfrak{B} r_{m+1} \\ = l_4(\mathbf{u}_m, \mathbf{h}_m, \rho_m) - l_4(\mathbf{u}_{m-1}, \mathbf{h}_{m-1}, \rho_{m-1}) + l_5(\rho_m) - l_5(\rho_{m-1}), \\ r_{m+1,t} + \mathbf{V}(x) \cdot \nabla_{\tau} r_{m+1} - \mathbf{w}_{m+1} \cdot \mathbf{N}(y) = l_6(\mathbf{u}_m, \rho_m) - l_6(\mathbf{u}_{m-1}, \rho_{m-1}), \quad y \in \mathcal{G}, \\ \mathbf{w}_{m+1}(y, 0) = 0, \quad y \in \mathcal{F}_1, \quad r_{m+1}(y, 0) = 0, \quad y \in \mathcal{G}, \end{array} \right. \quad (3.17)$$

$$\left\{ \begin{array}{l} \mu \mathbf{k}_{m+1,t} + \alpha^{-1} \text{rot} \text{rot} \mathbf{k}_{m+1} = \mathbf{l}_7(\mathbf{h}_m, \mathbf{u}_m, \rho_m) - \mathbf{l}_7(\mathbf{h}_{m-1}, \mathbf{u}_{m-1}, \rho_{m-1}), \\ \nabla \cdot \mathbf{k}_{m+1} = 0, \quad y \in \mathcal{F}_1, \\ \mu \mathbf{k}_{m+1,t} + \alpha^{-1} \text{rot} \text{rot} \mathbf{k}_{m+1} = 0, \quad \nabla \cdot \mathbf{k}_{m+1} = 0, \quad y \in \mathcal{F}_3, \\ \text{rot} \mathbf{k}_{m+1} = \text{rot}(\mathbf{l}_8(\mathbf{h}_m, \rho_m) - \mathbf{l}_8(\mathbf{h}_{m-1}, \rho_{m-1})), \quad \nabla \cdot \mathbf{k}_{m+1} = 0, \quad y \in \mathcal{F}_2, \\ [\mu \mathbf{k}_{m+1} \cdot \mathbf{N}] = 0, \quad [\mathbf{k}_{m+1, \tau}] = \mathbf{l}_9(\mathbf{h}_m, \rho_m) - \mathbf{l}_9(\mathbf{h}_{m-1}, \rho_{m-1}), \quad y \in \mathcal{G}, \\ [\mu \mathbf{k}_{m+1} \cdot \mathbf{n}] = 0, \quad [\mathbf{k}_{m+1, \tau}] = 0, \quad y \in S_3, \\ \mathbf{k}_{m+1} \cdot \mathbf{n} = 0, \quad y \in S, \\ \int_{\mathcal{F}_2} \mu \mathbf{k}_{m+1,t} \cdot \mathbf{u}_q^{\mathcal{F}} dy - \int_{\mathcal{G}} \mathbf{N} \times \alpha^{-1} \text{rot} \mathbf{k}_{m+1} \cdot \mathbf{u}_q^{\mathcal{F}} dS - \int_{S_3} \mathbf{n} \times \alpha^{-1} \text{rot} \mathbf{k}_{m+1} \cdot \mathbf{u}_q^{\mathcal{F}} dS \\ = l_{10,q}(\mathbf{u}_m, \rho_m, \mathbf{h}_m) - l_{10,q}(\mathbf{u}_{m-1}, \rho_{m-1}, \mathbf{h}_{m-1}), \quad q = 1, \dots, b_1(\mathcal{F}_2), \\ \mathbf{k}_{m+1}(y, 0) = 0, \quad y \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3. \end{array} \right. \quad (3.18)$$

Using Theorems 2, 4 and the estimates of the differences of nonlinear terms in (3.17), (3.18), we show that

$$Y(\mathbf{w}_{m+1}, s_{m+1}, r_{m+1}, \mathbf{k}_{m+1}) \leq \vartheta_1 Y(\mathbf{w}_m, s_m, r_m, \mathbf{k}_m),$$

$m > 1$ , with a small  $\vartheta_1$ , which guarantees the boundedness of

$$\sum_{m=2}^{\infty} Y(\mathbf{u}_{m+1} - \mathbf{u}_m, q_{m+1} - q_m, \rho_{m+1} - \rho_m, \mathbf{h}_{m+1} - \mathbf{h}_m)$$

and the convergence of  $(\mathbf{u}_m, q_m, \rho_m, \mathbf{h}_m)$ . The idea of the proof is the same as in [2, 7] in the case  $p, r = 2$ , and we omit the details.

It is clear that  $(\mathbf{u}_m, q_m, \rho_m, \mathbf{h}_m)$  tend to the solution of the problem (5.1), as  $m \rightarrow \infty$ , and the solution satisfies (3.14):

$$Y(T) \leq 2c_1 N. \quad (3.19)$$

## 4 Construction of the electric field

We have proved the existence of  $(\mathbf{u}, q, \rho, \mathbf{h})$  satisfying (1.6), (1.15), (1.16) and the additional condition (1.18). Now we find  $\mathbf{e}$  satisfying (1.7), (1.8). We introduce the vector field  $\mathbf{e}_1$  such that

$$\begin{aligned} \nabla \cdot \mathbf{e}_1(y, t) &= 0, \quad y \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3, \quad [\mathbf{N} \times \mathcal{P}\mathbf{e}_1] = \mathbf{\Psi}, \quad [\mathbf{N} \cdot \mathbf{e}_1] = 0, \quad y \in \mathcal{G}, \\ [\mathbf{n} \cdot \mathbf{e}_1] &= 0, \quad y \in S_3, \quad \mathbf{e}_1(y, t) = 0, \quad y \in S, \end{aligned} \quad (4.1)$$

$$\sum_{i=1}^3 \|\mathbf{e}_1\|_{W_r^{1,0}(Q_T^i)} \leq c \|\mathbf{\Psi}\|_{W_r^{1-1/r}(G_T)}. \quad (4.2)$$

Since the matrix  $\mathcal{P}$  is close to  $I$ , such  $\mathbf{e}_1$  exists. As shown in [1, Proposition 1], see also [3], the conditions

$$\nabla \cdot \mathbf{h}(y, t) = 0, \quad y \in \mathcal{F}_1 \cup \mathcal{F}_2, \quad [\mu \mathbf{h} \cdot \mathbf{N}] = 0, \quad [\mathbf{N} \times \mathcal{P}\mathbf{e}_1] = \mathbf{\Psi}, \quad y \in \mathcal{G}, \quad (4.3)$$

imply  $[\mathbf{N} \cdot \text{rot} \mathcal{P}\mathbf{e}_1] = -[\mu(\mathbf{h}_t - \mathbf{\Phi}) \cdot \mathbf{N}] = [\mu \mathbf{\Phi} \cdot \mathbf{N}]$ ,  $y \in \mathcal{G}$ . In addition, we have

$$\mathbf{n} \cdot ((\mu \mathbf{h}_t - \mathbf{\Phi}) + \text{rot} \mathcal{P}\mathbf{e}_1)|_{y \in S} = 0. \quad (4.4)$$

Now, we consider the problem

$$\text{rot} \mathcal{E} = -\mu(\mathbf{h}_t - \mathbf{\Phi}) - \text{rot} \mathcal{P}\mathbf{e}_1, \quad \nabla \cdot \mathcal{P}^{-1} \mathcal{E} = 0, \quad y \in \Omega, \quad \mathcal{E}_\tau = 0, \quad y \in S. \quad (4.5)$$

**Theorem 6.** *The problem (4.5) has a unique solution  $\mathcal{E} \in W_r^{1,0}(Q_T)$  orthogonal to the  $b_2(\Omega)$ -dimensional space  $\widetilde{U}_d(\Omega)$  of vector fields  $\mathbf{v}_j(y) = \nabla W_j(y)$  such that*

$$\nabla \cdot \mathcal{P}^{-1} \nabla W_j(y) = 0, \quad y \in \Omega, \quad W_j|_{S'_k} = \delta_{jk}, \quad W_j|_{S'_0} = 0, \quad (4.6)$$

where  $S'_k$ ,  $k = 0, \dots, b_2(\Omega)$ , are all the components of the boundary  $S$  of  $\Omega$ . The solution satisfies the inequality

$$\|\mathcal{E}\|_{W_r^{1,0}(Q_T)} \leq c(\|\mathbf{h}_t\|_{L_r(Q_T)} + \|\mathbf{\Phi}\|_{L_r(Q_T)} + \|\mathbf{\Psi}\|_{W_r^{1-1/r}(G_T)}). \quad (4.7)$$

**Proof.** As in [3], the solution may be expressed by the explicit formula

$$\begin{aligned} \mathcal{E}(y, t) &= \mathcal{E}_1(y, t) + \nabla Z(y, t) + \sum_{j=1}^{b_2(\Omega)} A_j(t) \nabla V_j(y), \\ \mathcal{E}_1(y, t) &= -\frac{1}{4\pi} \text{rot} \int_{\Omega} \frac{\mu(\mathbf{h}_t(z, t) - \mathbf{\Phi}(z, t)) + \text{rot} \mathcal{P}\mathbf{e}_1(z, t)}{|y - z|} dz, \\ \nabla \cdot \mathcal{P}^{-1} \nabla Z(y, t) &= -\nabla \cdot \mathcal{P}^{-1} \mathcal{E}_1, \quad y \in \Omega, \quad Z(y, t) = -g(y, t), \quad y \in S, \end{aligned}$$

The function  $g$  is defined as follows. In view of (4.4),  $\int_{\Sigma} \text{rot} \mathcal{E}_1 \cdot \mathbf{n} dS = 0$  for arbitrary  $\Sigma \subset S$  and, moreover, for the surfaces  $\Sigma_k \subset \Omega$  such that the domain  $\Omega \setminus \cup_k \Sigma_k$  is simply connected. This follows from the condition (1.18), i.e.,

$$\int_{\Omega} (\mu(\mathbf{h}_t - \mathbf{\Phi}) + \text{rot} \mathcal{P}\mathbf{e}_1) \cdot \mathbf{u}_q^\Omega dy = 0, \quad q = 1, \dots, b_1(\Omega),$$

see [8,9] for details. Hence  $\int_L \mathcal{E}_1 \cdot d\mathbf{l} = 0$  for arbitrary closed contour  $L \subset S$ , which means that  $\mathcal{E}_1 = \nabla_\tau g$  and  $\mathcal{E}_{1,\tau} + \nabla_\tau Z|_S = 0$ .

From the Calderon-Zygmund theorem and elliptic estimates it follows that

$$\|\mathcal{E}_1 + \nabla Z\|_{W_r^{1,0}(Q_T)} \leq c(\|\mathbf{h}_t\|_{L_r(Q_T)} + \|\Phi\|_{L_r(Q_T)} + \|\Psi\|_{W_r^{1-1/r,0}(G_T)}).$$

Moreover, the orthogonality condition

$$0 = \int_{\Omega} (\mathcal{E}_1 + \nabla Z + \sum_{j=1}^{b_2(\Omega)} A_k(t) \nabla V_j) \cdot \nabla V_q dy$$

yields the estimate of  $A_k$  by  $\mathcal{E}_1 + \nabla Z$ , which completes the proof of (4.7) and of Theorem 6.

Now we pass to the definition of  $\mathbf{e}$ . We set

$$\begin{aligned} \mathcal{P}\mathbf{e}^{(1)} &= \alpha_1^{-1} \mathcal{P} \text{rot} \mathcal{P}\mathbf{h} - \mu(\mathcal{L}^{-1} \mathbf{u} \times \mathbf{h}), \quad y \in \mathcal{F}_1, \\ \mathbf{e}^{(3)} &= \alpha^{-1}(\text{rot} \mathbf{h} - \mathbf{j}), \quad y \in \mathcal{F}_3. \end{aligned} \quad (4.8)$$

Then

$$\begin{aligned} \text{rot} \mathcal{P}\mathbf{e} &= -\mu \mathbf{h}_t + \mu \Phi = \text{rot} \mathcal{E} + \text{rot} \mathcal{P}\mathbf{e}_1, \quad y \in \mathcal{F}_1, \\ \text{rote} &= \text{rot} \mathcal{E} + \text{rote}_1, \quad y \in \mathcal{F}_3, \end{aligned} \quad (4.9)$$

which implies

$$\begin{aligned} \mathcal{P}\mathbf{e} &= \mathcal{E} + \mathcal{P}\mathbf{e}_1 + \nabla \chi_1, \quad y \in \mathcal{F}_1, \\ \mathbf{e} &= \mathcal{E} + \mathbf{e}_1 + \nabla \chi_3, \quad y \in \mathcal{F}_3. \end{aligned} \quad (4.10)$$

Since  $\mathcal{F}_1$  and  $\mathcal{F}_3$  may be multi-connected, this conclusion should be justified. We show that

$$\int_{\mathcal{F}_1} (\mathcal{P}\mathbf{e} - \mathcal{E} - \mathcal{P}\mathbf{e}_1) \cdot \text{rot} \psi(y) dy + \int_{\mathcal{F}_3} (\mathbf{e} - \mathcal{E} - \mathbf{e}_1) \cdot \text{rot} \psi(y) dy = 0, \quad \forall \psi \in \mathcal{H}^{(1)}(\Omega), \quad (4.11)$$

where  $\mathcal{H}^{(1)}(\Omega)$  is the space of solenoidal vector fields from  $W_2^1(\mathcal{F}_1) \cap W_2^1(\mathcal{F}_2) \cap W_2^1(\mathcal{F}_3)$  satisfying the equations  $\text{rot} \psi(y) = 0$ ,  $\nabla \cdot \psi(y) = 0$  in  $\mathcal{F}_2$  and the relations (2.16), (2.17) with  $k_i = \text{const}$  (cf. [12], [9]). Then (4.10) follows from Theorem 3 in [2] (with  $\mathbf{a} = 0$ ,  $\mathbf{j}(y) = 0$  in  $\mathcal{F}_2$ ), that shows that  $\text{rot} \psi|_{\mathcal{F}_i}$ ,  $i = 1, 3$ , coincides with the space of solenoidal vector fields in  $\mathcal{F}_i$  with vanishing normal components on the boundaries  $\mathcal{G}$  and  $S_3$ ; see also Theorem 3.3 in [9].

Let  $\mathcal{H}_0^{(1)}(\Omega)$  be the subspace of  $\mathcal{H}^{(1)}(\Omega)$  such that  $\psi(y) = \nabla \varphi(y)$ ,  $y \in \mathcal{F}_2$  and  $k_i = 0$  in (2.16). The remaining finite dimensional subspace of  $\mathcal{H}^{(1)}(\Omega)$  is a linear combination of vector fields  $\psi_q(y)$ ,  $q = 1, \dots, b_1(\mathcal{F}_2)$  such that

$$\begin{aligned} \psi_q(y) &= \mathbf{u}_q^{\mathcal{F}}(y), \quad \mathbf{u}_q^{\mathcal{F}} \in U_n(\mathcal{F}_2), \quad y \in \mathcal{F}_2, \\ \nabla \cdot \psi_q(y) &= 0, \quad y \in \mathcal{F}_1, \quad \psi_q(y) \cdot \mathbf{N}(y) = 0, \quad (\psi_q)_\tau = (\mathbf{u}_q)_\tau(y), \quad y \in \mathcal{G}, \\ \nabla \cdot \psi_q(y) &= 0, \quad y \in \mathcal{F}_3, \quad \psi_q(y) \cdot \mathbf{n}(y) = 0, \quad (\psi_q)_\tau = (\mathbf{u}_q)_\tau(y), \quad y \in S_3. \end{aligned}$$

Let  $\psi \in \mathcal{H}_0^{(1)}(\Omega)$  and let  $\Phi(y)$ ,  $y \in \mathcal{F}_1 \cup \mathcal{F}_3$  be the solution of

$$\begin{aligned} \nabla^2 \Phi(y) &= 0, \quad y \in \mathcal{F}_1, \quad \Phi(y) = \varphi(y), \quad y \in \mathcal{G}, \\ \nabla^2 \Phi(y) &= 0, \quad y \in \mathcal{F}_3, \quad \Phi(y) = \varphi(y), \quad y \in S_3. \end{aligned}$$

Since  $(\varphi(y) - \nabla\Phi(y))_\tau = 0$ ,  $y \in \mathcal{G} \cup S_3$ , we have

$$\begin{aligned}
& \int_{\mathcal{F}_1} (\mathcal{P}\mathbf{e} - \mathcal{E} - \mathcal{P}\mathbf{e}_1) \cdot \text{rot}\boldsymbol{\psi}(y)dy + \int_{\mathcal{F}_3} (\mathbf{e} - \mathcal{E} - \mathbf{e}_1) \cdot \text{rot}\boldsymbol{\psi}(y)dy \\
&= \int_{\mathcal{F}_1} (\mathcal{P}\mathbf{e} - \mathcal{E} - \mathcal{P}\mathbf{e}_1) \cdot \text{rot}(\boldsymbol{\psi}(y) - \nabla\Phi(y))dy + \int_{\mathcal{F}_3} (\mathbf{e} - \mathcal{E} - \mathbf{e}_1) \cdot \text{rot}(\boldsymbol{\psi}(y) - \nabla\Phi(y))dy \\
&= \int_{\mathcal{F}_1} \text{rot}(\mathcal{P}\mathbf{e} - \mathcal{E} - \mathcal{P}\mathbf{e}_1) \cdot (\boldsymbol{\psi}(y) - \nabla\Phi(y))dy \\
&+ \int_{\mathcal{F}_3} \text{rot}(\mathbf{e} - \mathcal{E} - \mathbf{e}_1) \cdot (\boldsymbol{\psi}(y) - \nabla\Phi(y))dy = 0,
\end{aligned} \tag{4.12}$$

$\forall \boldsymbol{\psi} \in \mathcal{H}_0^{(1)}(\Omega)$ . In addition, if  $\boldsymbol{\psi} = \boldsymbol{\psi}_q$ , then

$$\begin{aligned}
& \int_{\mathcal{F}_1} (\mathcal{P}\mathbf{e} - \mathcal{E} - \mathcal{P}\mathbf{e}_1) \cdot \text{rot}\boldsymbol{\psi}_q(y)dy + \int_{\mathcal{F}_3} (\mathbf{e} - \mathcal{E} - \mathbf{e}_1) \cdot \text{rot}\boldsymbol{\psi}_q(y)dy \\
&= \int_{\mathcal{F}_1} (\alpha^{-1}\mathcal{P}\text{rot}\mathcal{P}\mathbf{h} - \mathbf{J}) \cdot \text{rot}\boldsymbol{\psi}_q(y)dy + \int_{\mathcal{F}_3} \alpha^{-1}(\text{rot}\mathbf{h} - \mathbf{j}) \cdot \text{rot}\boldsymbol{\psi}_q(y)dy \\
&+ \int_{\mathcal{F}_1 \cup \mathcal{F}_3} \mu(\mathbf{h}_t - \boldsymbol{\Phi}) \cdot \boldsymbol{\psi}_q dy \\
&+ \int_{\mathcal{G}} \mathbf{N} \times (\mathcal{E} + \mathcal{P}\mathbf{e}_1^{(1)}) \cdot \boldsymbol{\psi}_q(y)dS + \int_{S_3} \mathbf{n} \times (\mathcal{E} + \mathbf{e}_1^{(3)}) \cdot \boldsymbol{\psi}_q(y)dS,
\end{aligned} \tag{4.13}$$

and since the last two surface integrals are equal to

$$- \int_{\mathcal{F}_2} \text{rot}(\mathcal{E} + \mathcal{P}\mathbf{e}_1) \cdot \boldsymbol{\psi}_q(y)dy = \int_{\mathcal{F}_2} \mu(\mathbf{h}_t - \boldsymbol{\Phi}) \cdot \boldsymbol{\psi}_q dy + \int_{\mathcal{G}} \boldsymbol{\Psi} \cdot \boldsymbol{\psi}_q dS,$$

we can conclude from (4.10) and (1.16) that

$$\int_{\mathcal{F}_1} (\mathcal{P}\mathbf{e} - \mathcal{E} - \mathcal{P}\mathbf{e}_1) \cdot \text{rot}\boldsymbol{\psi}_q(y)dy + \int_{\mathcal{F}_3} (\mathbf{e} - \mathcal{E} - \mathbf{e}_1) \cdot \text{rot}\boldsymbol{\psi}_q(y)dy = 0,$$

q.e.d.

Finally, we set

$$\mathcal{P}\mathbf{e} = \mathcal{E} + \mathcal{P}\mathbf{e}_1 + \nabla\chi_2 + \sum_{j=1}^{b_2(\mathcal{F}_2)} D_j(t)\nabla W_j(y), \quad y \in \mathcal{F}_2, \tag{4.14}$$

where  $W_j$  and  $\chi_2$  are solutions to the problems (4.6) and

$$\begin{aligned}
& \nabla \cdot \mathcal{P}^{-1}\nabla\chi_2(y, t) = 0, \quad y \in \mathcal{F}_2, \\
& \chi_2(y, t) = \chi_1(y, t), \quad y \in \mathcal{G}, \quad \chi_2(y, t) = \chi_3(y, t), \\
& y \in S_3, \quad \chi_2(y, t) = 0, \quad y \in S.
\end{aligned} \tag{4.15}$$

It is easily verified that the vector field  $\mathbf{e}$  defined by (4.10), (4.14) satisfies (1.7), (1.8). In order to find  $D_j(t)$ , we need to impose on  $\mathbf{e}^{(2)}$  some normalization restrictions, for instance of the form

$$\int_{S'_k} \mathcal{P}\mathbf{e}^{(2)}(y, t) \cdot \mathbf{n}(y)dS = \int_{S'_k} \mathbf{e}^{(2)}(y, t) \cdot \mathbf{n}(y)dS = 0, \quad k = 1, \dots, b_2(\mathcal{F}_2). \tag{4.16}$$

Since the matrix  $\mathcal{J}$  with the elements  $\int_{S_k} \frac{\partial W_i}{\partial n} dS$ ,  $j, k = 1, \dots, b_2(\mathcal{F}_2)$  is non-degenerate, the equations (4.14), (4.16) define  $D_j$  in a unique way.

As in [3], the functions  $\mathbf{e}$ ,  $\nabla \chi_i$ ,  $D_j$  are easily estimated with the help of the relations (4.8), (4.14), and these estimates show that  $\mathbf{e} \in W_r^{1,0}(Q_T^i)$ .

But if  $\mathbf{h}_0 \in W_p^{2-2/p}(\mathcal{F}_i)$ ,  $\mathbf{j} \in W_p^{1,0}(\mathcal{F}_2)$ , then  $\mathbf{h} \in W_p^{2,1}(Q_T^i)$ ,  $\mathbf{e} \in W_p^{1,0}(Q_T^i)$ ,  $i = 1, 2, 3$  (cf. the remark at the end of Sec. 5 in [3]).

## References

1. V.A.Solonnikov, *Free boundary problems of magnetohydrodynamics in multi-connected domains*, Interfaces and free boundaries, **14** (2012), 569-603.
2. M.Padula, V.A.Solonnikov, *On the free boundary problem of magnetohydrodynamics*, J. Math. Sci. **178** No 3 (2011), 313-344.
3. V.A.Solonnikov,  *$L_p$ -theory of free boundary problems of magnetohydrodynamics in simply connected domains*, submitted to Trudy Spb. Math. Soc.
4. L.D.Landau, E.M. Lifshitz, *Electrodynamics of continuous media*, Pergamon Press (1960).
5. P.H.Roberts, *An introduction in magnetohydrodynamics*, London, Longmans, Green and Co. LTD (1967).
6. O.V.Besov, V.P.Il'in, S.M.Nikol'skii, *Integral representations of functions and imbedding theorems*, Moscow, "Nauka", 1975.
7. V.A.Solonnikov, E.V.Frolova,  *$L_p$ -theory for the Stefan problem*, Zapiski Nauchn. Semin. POMI **243** (1997), 299-323.
8. E. B. Bykhovskii, N. V. Smirnov, *On the orthogonal decomposition of the space of vector functions square summable in a given domain and the operators of vector analysis*, Tr. Mat. Inst. Steklov, **59** (1960), 6-36.
9. S. Mosconi, V.A.Solonnikov, *On a problem of magnetohydrodynamics in a multi-connected domain*, Nonlinear Analysis **74**, No. 2 (2010), 462-478.
10. V.A.Solonnikov,  *$L_p$ -estimates for the linear problem arising in the study of the motion of an isolated liquid mass*. J.Math. Sci. 189, No 4 (2013), p. 699.
11. V.A.Solonnikov,  *$L_p$  -estimates for the solution of a linear problem that arises in magneto-hydrodynamics. (Russian)* Algebra i Analiz **23** (2011), no. 1, 232-254.
12. O. A. Ladyzhenskaya, V. A. Solonnikov. *The linearization principle and invariant manifolds for problems of magnetohydrodynamics*, J. of Math. Sci., **8** (1977), 384-422.