

P. Neff, D. Pauly, K.-J. Witsch

ON A CANONICAL EXTENSION OF KORN'S FIRST
AND POINCARÉ'S INEQUALITY TO $H(\text{CURL})$

ABSTRACT. We prove a Korn-type inequality in $\mathring{H}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$ for tensor fields P mapping Ω to $\mathbb{R}^{3 \times 3}$. More precisely, let $\Omega \subset \mathbb{R}^3$ be a bounded domain with connected Lipschitz boundary $\partial\Omega$. Then, there exists a constant $c > 0$ such that

$$c \|P\|_{L^2(\Omega, \mathbb{R}^{3 \times 3})} \leq \|\text{sym } P\|_{L^2(\Omega, \mathbb{R}^{3 \times 3})} + \|\text{Curl } P\|_{L^2(\Omega, \mathbb{R}^{3 \times 3})} \quad (0.1)$$

holds for all tensor fields $P \in \mathring{H}(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$, i.e., all

$$P \in H(\text{Curl}; \Omega, \mathbb{R}^{3 \times 3})$$

with vanishing tangential trace on $\partial\Omega$. Here, rotation and tangential trace are defined row-wise. For compatible P (i.e., $P = \nabla v$), $\text{Curl } P = 0$, where $v \in H^1(\Omega, \mathbb{R}^3)$ a vector field having components v_n , for which ∇v_n are normal at $\partial\Omega$, the estimate (0.1) is reduced to a non-standard variant of the Korn's first inequality:

$$c \|\nabla v\|_{L^2(\Omega, \mathbb{R}^{3 \times 3})} \leq \|\text{sym } \nabla v\|_{L^2(\Omega, \mathbb{R}^{3 \times 3})}.$$

For skew-symmetric P ($\text{sym } P = 0$) the estimate (0.1) generates a non-standard version of the Poincaré. Therefore, the estimate is a generalization of two classical inequalities of Poincaré and Korn.

§1. INTRODUCTION: INFINITESIMAL GRADIENT PLASTICITY

The motivation for our new estimate is a formulation of infinitesimal gradient plasticity [2]. Our model is taken from Neff et al. [9]. Let $\Omega \subset \mathbb{R}^3$ be a bounded domain. The goal is to find the displacement $u : [0, \infty) \times \Omega \mapsto \mathbb{R}^3$ and the possibly non-symmetric plastic distortion tensor $P : [0, \infty) \times \Omega \mapsto \mathbb{R}^{3 \times 3}$, such that in $[0, \infty) \times \Omega$

$$\text{Div } \sigma = f, \quad \sigma = 2\mu \text{sym}(\nabla u - P) + \lambda \text{tr}(\nabla u - P) \text{id}, \quad (1.1)$$

$$\dot{P} \in \Phi(\Sigma), \quad \Sigma = \sigma - 2\mu \text{sym } P - \mu L_c^2 \text{Curl } \text{Curl } P, \quad (1.2)$$

hold. The system is completed by the boundary conditions

$$u(t, x) = 0, \quad \nu(x) \times P(t, x) = 0 \quad \forall (t, x) \in [0, \infty) \times \partial\Omega \quad (1.3)$$

Key words and phrases: Korn's inequality, gradient plasticity, theory of Maxwell's equations, Helmholtz' decomposition, Poincaré/Friedrichs type estimate.

and the initial condition $P(0, x) = 0$ for all $x \in \Omega$. The underlying thermodynamic potential including the plastic gradients in form of the dislocation density tensor $\text{Curl } P$ is

$$\int_{\Omega} \mu |\text{sym}(\nabla u - P)|^2 + \frac{\lambda}{2} |\text{tr}(\nabla u - P)|^2 - f \cdot u + \mu |\text{sym } P|^2 + \frac{\mu}{2} L_c^2 |\text{Curl } P|^2.$$

Here, μ, λ are the elastic Lamé moduli and σ is the symmetric Cauchy stress tensor. The system is driven by nonzero body forces denoted by f . The exterior normal to the boundary $\partial\Omega$ is denoted by ν and the plastic distortion P is required to satisfy row-wise the homogeneous tangential boundary condition which means that the boundary $\partial\Omega$ is a perfect conductor regarding the plastic distortion.¹

Moreover, $\Phi : \mathbb{R}^{3 \times 3} \mapsto \mathbb{R}^{3 \times 3}$ is the monotone, multivalued flow-function with $\Phi(0) = 0$ and $\Phi(\mathbb{R}_{\text{sym}}^{3 \times 3}) \subset \mathbb{R}_{\text{sym}}^{3 \times 3}$. In general, Σ is not symmetric even if P is symmetric. Thus, the plastic inhomogeneity is responsible for the plastic spin (the possible non-symmetry of P). The mathematically suitable space for symmetric plastic distortion P is the classical space $\mathbf{H}(\text{curl}; \Omega)$ for each row of P [2, 13]. This case appears when choosing $\Phi : \mathbb{R}^{3 \times 3} \mapsto \mathbb{R}_{\text{sym}}^{3 \times 3}$.

In the large scale limit $L_c \rightarrow 0$ we recover a classical elasto-plasticity model with local kinematic hardening and symmetric plastic strain $\varepsilon_P := \text{sym } P$, since then $\dot{P} \in \mathbb{R}_{\text{sym}}^{3 \times 3}$.

Uniqueness of classical solutions for rate-independent and rate-dependent formulations of this model is shown in [9]. The more difficult existence question for the rate-independent model in terms of a weak reformulation is addressed in [9]. First numerical results for a simplified rate-independent irrotational formulation (no plastic spin, i.e., symmetric plastic distortion P) are presented in [13], cf [19]. In [3] the model has been extended to rate-independent isotropic hardening based on the concept of a dissipation function defined in terms of the equivalent plastic strain. From a modeling point of view, it is strongly preferable to again have only the symmetric (rate) part of the plastic distortion P appear in the dissipation potential.

The existence and uniqueness can be settled by recasting the model as a variational inequality, if it is possible to define a bilinear form which

¹This homogeneous tangential boundary condition on P is consistent with $\nu \times \nabla u = 0$ on $\partial\Omega$ which follows from $u = 0$ on $\partial\Omega$.

is coercive with respect to appropriate spaces. This program has been achieved for other variants of the model in [3]. It had to remain basically open for the above system (1.1)–(1.3). In this case, the appropriate space for the plastic distortion P is the completion $\mathring{H}_{\text{sym}}(\text{Curl}; \Omega)$ of the linear space

$$\{P \in C^\infty(\bar{\Omega}, \mathbb{R}^{3 \times 3}) : P_n \text{ normal at } \partial\Omega, n = 1, 2, 3\}$$

with respect to the norm $\|\cdot\|$, where P_n are the columns of P^t and

$$\|P\|^2 := \|\text{sym } P\|_{L^2(\Omega)}^2 + \|\text{Curl } P\|_{L^2(\Omega)}^2.$$

Despite first appearance, this quadratic form indeed defines a norm on $\mathring{H}_{\text{sym}}(\text{Curl}; \Omega)$ since for skew-symmetric tensors the operator Curl controls all derivatives, i.e., the full gradient. This was already mentioned in [9]. Thus $\mathring{H}_{\text{sym}}(\text{Curl}; \Omega)$ is a Hilbert-space. However, it is not clear that this space admits a (linear and bounded) tangential trace operator. Since only $\|\text{sym } P\|_{L^2(\Omega)}$ appears in $\|P\|$, it is also not clear at all, if the skew-symmetric part of P is controlled. Therefore, the crucial embedding

$$\mathring{H}_{\text{sym}}(\text{Curl}; \Omega) \subset L^2(\Omega)$$

might fail. As a consequence of our main result of this paper we obtain that nevertheless

$$\mathring{H}_{\text{sym}}(\text{Curl}; \Omega) = \mathring{H}(\text{Curl}; \Omega)$$

holds with equivalent norms in case the domain Ω has a connected Lipschitz boundary.

The result of this paper has been announced in [10,11] and a forthcoming paper [12] will be devoted to the case $\Omega \subset \mathbb{R}^N$.

For the proof of our main result (0.1) we combine techniques from electro-magnetic and elastic theory, namely the Helmholtz decomposition, the Maxwell compactness property and Korn's first inequality. Their basic variants are well known results which can be found in many books, e.g., in [6] and the literature cited there. More sophisticated and related versions are presented, e.g., in [14, 16–18, 23] for Maxwell's equations and [1, 8] for Korn's inequality.

The paper at hand is organized as follows: After this preliminary motivation we introduce our notations, definitions and provide some background results. In section 3 we give the proof of our main estimate. In the last

section 4 we establish a connection to a related result by Garroni et al. [4] for the two-dimensional case.

§2. DEFINITIONS AND PRELIMINARIES

Let Ω be a bounded domain in \mathbb{R}^3 with connected Lipschitz continuous boundary $\Gamma := \partial\Omega$.

2.1. Functions and vector fields. The usual Lebesgue spaces of square integrable functions, vector or tensor fields on Ω with values in \mathbb{R} , \mathbb{R}^3 or $\mathbb{R}^{3 \times 3}$, respectively, will be denoted by $L^2(\Omega)$. Moreover, we introduce the standard Sobolev spaces

$$\begin{aligned} \mathbf{H}(\text{grad}; \Omega) &= \{u \in L^2(\Omega) : \text{grad } u \in L^2(\Omega)\}, \\ \|u\|_{\mathbf{H}(\text{Grad}; \Omega)}^2 &:= \|u\|_{L^2(\Omega)}^2 + \|\text{grad } u\|_{L^2(\Omega)}^2, \\ \mathbf{H}(\text{curl}; \Omega) &= \{v \in L^2(\Omega) : \text{curl } v \in L^2(\Omega)\}, \\ \|v\|_{\mathbf{H}(\text{curl}; \Omega)}^2 &:= \|v\|_{L^2(\Omega)}^2 + \|\text{curl } v\|_{L^2(\Omega)}^2, \\ \mathbf{H}(\text{div}; \Omega) &= \{v \in L^2(\Omega) : \text{div } v \in L^2(\Omega)\}, \\ \|v\|_{\mathbf{H}(\text{div}; \Omega)}^2 &:= \|v\|_{L^2(\Omega)}^2 + \|\text{div } v\|_{L^2(\Omega)}^2. \end{aligned}$$

$\mathbf{H}(\text{grad}; \Omega)$ is often denoted by $\mathbf{H}^1(\Omega)$. Furthermore, we define their closed subspaces $\mathring{\mathbf{H}}(\text{grad}; \Omega)$, $\mathring{\mathbf{H}}(\text{curl}; \Omega)$ as completion under the respective norms of the scalar resp. vector valued space $\mathring{\mathbf{C}}^\infty(\Omega)$ of compactly supported and smooth test functions resp. vector fields. In the latter Sobolev spaces the usual homogeneous scalar resp. tangential boundary conditions

$$u|_\Gamma = 0, \quad \nu \times v|_\Gamma = 0$$

are generalized, where ν denotes the outer unit normal at Γ . We note in passing that $\nu \times v|_\Gamma = 0$ is equivalent to $\tau \cdot v|_\Gamma = 0$ for all tangential directions τ at Γ , which means that v is normal at Γ . Furthermore, we need the spaces of irrotational or solenoidal vector fields

$$\begin{aligned} \mathbf{H}(\text{curl}_0; \Omega) &:= \{v \in \mathbf{H}(\text{curl}; \Omega) : \text{curl } v = 0\}, \\ \mathring{\mathbf{H}}(\text{curl}_0; \Omega) &:= \{v \in \mathring{\mathbf{H}}(\text{curl}; \Omega) : \text{curl } v = 0\}, \\ \mathbf{H}(\text{div}_0; \Omega) &:= \{v \in \mathbf{H}(\text{div}; \Omega) : \text{div } v = 0\}, \end{aligned}$$

where the index 0 indicates vanishing curl or div, respectively. All these spaces are Hilbert spaces. E.g., in classical terms we have $v \in \mathring{H}(\text{curl}_0; \Omega)$, if and only if

$$\text{curl } v = 0, \quad \nu \times v|_{\Gamma} = 0.$$

For an introduction of these spaces see [6, p. 11–12, 148] or [5, p. 26]. The most important tool for our analysis is the compact embedding

$$\mathring{H}(\text{curl}; \Omega) \cap H(\text{div}; \Omega) \hookrightarrow L^2(\Omega),$$

which is often referred as ‘Maxwell compactness property’, see [6, p. 158] and [16, 18, 21, 23, 24]. A first immediate consequence is that the space of so called ‘harmonic Dirichlet fields’

$$\mathcal{H}(\Omega) := \mathring{H}(\text{curl}_0; \Omega) \cap H(\text{div}_0; \Omega)$$

is finite dimensional. A vector field v belonging to $\mathcal{H}(\Omega)$ means in classical terms that

$$\text{curl } v = 0, \quad \text{div } v = 0, \quad \nu \times v|_{\Gamma} = 0.$$

The dimension of $\mathcal{H}(\Omega)$ equals the second Betti number of Ω , see [6, p. 159] and [15, Theorem 1]. Since we assume the boundary Γ to be connected, there are no Dirichlet fields besides zero, i.e.,

$$\mathcal{H}(\Omega) = \{0\}.$$

This condition on the domain Ω resp. its boundary Γ is satisfied e.g. for a ball or a torus.

By a usual indirect argument we achieve another immediate consequence, see [6, p. 158, Theorem 8.9] or [5, Lemma 3.4]:

Lemma 1. (Maxwell Estimate for Vector Fields) *There exists a positive constant c_m , such that for all vector fields $v \in \mathring{H}(\text{curl}; \Omega) \cap H(\text{div}; \Omega)$*

$$\|v\|_{L^2(\Omega)} \leq c_m \left(\|\text{curl } v\|_{L^2(\Omega)}^2 + \|\text{div } v\|_{L^2(\Omega)}^2 \right)^{1/2}.$$

By definition of the weak divergence, the projection theorem and Rellich’s selection theorem [6, p. 14] we have from [6, p. 148, Theorem 8.3] or [22, Lemma 3.5], [7, Theorem 3.45]

Lemma 2. (Helmholtz Decomposition for Vector Fields) *We have the orthogonal decomposition*

$$L^2(\Omega) = \text{grad } \mathring{H}(\text{grad}; \Omega) \oplus H(\text{div}_0; \Omega).$$

2.2. Tensor fields. We extend our calculus to (3×3) -tensor (matrix) fields. For vector fields v with components in $\mathbf{H}(\text{grad}; \Omega)$ and tensor fields T with rows in $\mathbf{H}(\text{curl}; \Omega)$ resp. $\mathbf{H}(\text{div}; \Omega)$, i.e.,

$$v = \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix}, \quad v_n \in \mathbf{H}(\text{grad}; \Omega),$$

$$T^t = [T_1 \ T_2 \ T_3], \quad T_n \in \mathbf{H}(\text{curl}; \Omega) \text{ resp. } \mathbf{H}(\text{div}; \Omega)$$

we define

$$\text{Grad } v := \begin{bmatrix} \text{grad}^t v_1 \\ \text{grad}^t v_2 \\ \text{grad}^t v_3 \end{bmatrix} = J_v = \nabla v, \quad \text{Curl } T := \begin{bmatrix} \text{curl}^t T_1 \\ \text{curl}^t T_2 \\ \text{curl}^t T_3 \end{bmatrix}, \quad \text{Div } T := \begin{bmatrix} \text{div } T_1 \\ \text{div } T_2 \\ \text{div } T_3 \end{bmatrix},$$

where J_v denotes the Jacobian of v and t the transpose. We note that v and $\text{Div } T$ are vector fields, whereas T , $\text{Curl } T$ and $\text{Grad } v$ are tensor fields. The corresponding Sobolev spaces will be denoted by $\mathbf{H}(\text{Grad}; \Omega)$, $\mathring{\mathbf{H}}(\text{Grad}; \Omega)$, $\mathbf{H}(\text{Curl}; \Omega)$, $\mathring{\mathbf{H}}(\text{Curl}; \Omega)$, $\mathbf{H}(\text{Curl}_0; \Omega)$, $\mathring{\mathbf{H}}(\text{Curl}_0; \Omega)$, $\mathbf{H}(\text{Div}; \Omega)$, $\mathbf{H}(\text{Div}_0; \Omega)$. As usual, we denote by $\text{sym } T := 1/2(T + T^t)$ the symmetric part of a tensor T .

Let us now present our three crucial tools to prove the new estimate. First we have obvious consequences from Lemmas 1 and 2:

Corollary 3. (Maxwell Estimate for Tensor Fields) *For all $T \in \mathring{\mathbf{H}}(\text{Curl}; \Omega) \cap \mathbf{H}(\text{Div}; \Omega)$*

$$\|T\|_{\mathbf{L}^2(\Omega)} \leq c_m (\|\text{Curl } T\|_{\mathbf{L}^2(\Omega)}^2 + \|\text{Div } T\|_{\mathbf{L}^2(\Omega)}^2)^{1/2}.$$

Corollary 4. (Helmholtz Decomposition for Tensor Fields) *We have the orthogonal decomposition*

$$\mathbf{L}^2(\Omega) = \text{Grad } \mathring{\mathbf{H}}(\text{Grad}; \Omega) \oplus \mathbf{H}(\text{Div}_0; \Omega).$$

The third important tool is Korn's first inequality [6, p. 207] or [20, p. 54]:

Lemma 5. (Korn's First Inequality) *For all $v \in \mathring{\mathbf{H}}(\text{Grad}; \Omega)$*

$$\|\text{Grad } v\|_{\mathbf{L}^2(\Omega)} \leq \sqrt{2} \|\text{sym Grad } v\|_{\mathbf{L}^2(\Omega)}.$$

§3. MAIN RESULTS

For tensor fields $T \in \mathbf{H}(\text{Curl}; \Omega)$ we define the semi-norm

$$\|T\| := \left(\|\text{sym } T\|_{\mathbf{L}^2(\Omega)}^2 + \|\text{Curl } T\|_{\mathbf{L}^2(\Omega)}^2 \right)^{1/2}.$$

Lemma 6. *Let $\hat{c} := \max\{2, \sqrt{5}c_m\}$. Then, for all $T \in \mathring{\mathbf{H}}(\text{Curl}; \Omega)$*

$$\|T\|_{\mathbf{L}^2(\Omega)} \leq \hat{c} \|T\|.$$

Proof. Let $T \in \mathring{\mathbf{H}}(\text{Curl}; \Omega)$. According to Corollary 4 we orthogonally decompose

$$T = \text{Grad } v + S \in \text{Grad } \mathring{\mathbf{H}}(\text{Grad}; \Omega) \oplus \mathbf{H}(\text{Div}_0; \Omega).$$

Then, $\text{Curl } T = \text{Curl } S$ and we observe $S \in \mathring{\mathbf{H}}(\text{Curl}; \Omega) \cap \mathbf{H}(\text{Div}_0; \Omega)$ since

$$\text{Grad } \mathring{\mathbf{H}}(\text{Grad}; \Omega) \subset \mathring{\mathbf{H}}(\text{Curl}_0; \Omega). \quad (3.1)$$

By Corollary 3 we have

$$\|S\|_{\mathbf{L}^2(\Omega)} \leq c_m \|\text{Curl } T\|_{\mathbf{L}^2(\Omega)}. \quad (3.2)$$

Then, by Lemma 5 and (3.2) we obtain easily

$$\begin{aligned} \|T\|_{\mathbf{L}^2(\Omega)}^2 &= \|\text{Grad } v\|_{\mathbf{L}^2(\Omega)}^2 + \|S\|_{\mathbf{L}^2(\Omega)}^2 \\ &\leq 2 \|\text{sym Grad } v\|_{\mathbf{L}^2(\Omega)}^2 + \|S\|_{\mathbf{L}^2(\Omega)}^2 \leq 4 \|\text{sym } T\|_{\mathbf{L}^2(\Omega)}^2 + 5 \|S\|_{\mathbf{L}^2(\Omega)}^2, \end{aligned}$$

which completes the proof. \square

An immediate consequence is

Theorem 7. *On $\mathring{\mathbf{H}}(\text{Curl}; \Omega)$ the norms $\|\cdot\|_{\mathbf{H}(\text{Curl}; \Omega)}$ and $\|\cdot\|$ are equivalent. In particular, $\|\cdot\|$ is a norm on $\mathring{\mathbf{H}}(\text{Curl}; \Omega)$ and*

$$\exists c > 0 \quad \forall T \in \mathring{\mathbf{H}}(\text{Curl}; \Omega) \quad c \|T\|_{\mathbf{H}(\text{Curl}; \Omega)} \leq \|\text{sym } T\|_{\mathbf{L}^2(\Omega)} + \|\text{Curl } T\|_{\mathbf{L}^2(\Omega)}.$$

3.1. Consequences for irrotational tensors: Korn's first inequality. Picking irrotational tensor fields T or setting $T := \text{Grad } v$, we obtain by Lemma 6 and (3.1)

Corollary 8. (Korn's First Inequality: Tangential-Variants)

$$(i) \quad \|T\|_{\mathbf{L}^2(\Omega)} \leq \hat{c} \|\text{sym } T\|_{\mathbf{L}^2(\Omega)} \text{ holds for all tensor fields } T \in \mathring{\mathbf{H}}(\text{Curl}_0; \Omega).$$

- (ii) $\|\text{Grad } v\|_{\mathbb{L}^2(\Omega)} \leq \widehat{c} \|\text{sym Grad } v\|_{\mathbb{L}^2(\Omega)}$ holds for all vector fields $v \in \mathring{\mathbb{H}}(\text{Grad}; \Omega)$ with $\text{Grad } v \in \mathring{\mathbb{H}}(\text{Curl}_0; \Omega)$.
- (iii) $\|\text{Grad } v\|_{\mathbb{L}^2(\Omega)} \leq \widehat{c} \|\text{sym Grad } v\|_{\mathbb{L}^2(\Omega)}$ holds for all vector fields $v \in \mathring{\mathbb{H}}(\text{Grad}; \Omega)$.

These are different but equivalent versions of Korn's first inequality from Lemma 5 with a larger constant \widehat{c} , since the boundary Γ is connected, i.e., $\mathcal{H}(\Omega) = \{0\}$, and hence

$$\text{Grad } \mathring{\mathbb{H}}(\text{Grad}; \Omega) = \mathring{\mathbb{H}}(\text{Curl}_0; \Omega).$$

Thus, e.g., (ii) holds, where the boundary condition means that $\text{Grad } v_n$, $n = 1, 2, 3$, are normal at Γ , which then extends Lemma 5 through the (apparently) weaker boundary condition.

3.2. Consequences for skew-symmetric tensors: Poincaré's inequality. Taking special skew-symmetric tensor fields

$$T = \begin{bmatrix} 0 & 0 & u \\ 0 & 0 & 0 \\ -u & 0 & 0 \end{bmatrix} \cong u, \quad \text{Curl } T = \begin{bmatrix} \partial_2 u & -\partial_1 u & 0 \\ 0 & 0 & 0 \\ 0 & -\partial_3 u & \partial_2 u \end{bmatrix}$$

with some scalar function u , we get by Lemma 6 since now $\text{Curl } T$ is as good as $\text{grad } u$ and by

$$\nu \times T|_{\partial\Omega} = \begin{bmatrix} \nu_2 u|_{\partial\Omega} & -\nu_1 u|_{\partial\Omega} & 0 \\ 0 & 0 & 0 \\ 0 & -\nu_3 u|_{\partial\Omega} & \nu_2 u|_{\partial\Omega} \end{bmatrix}$$

$\nu \times T|_{\partial\Omega} = 0$ is equivalent to $u|_{\partial\Omega} = 0$

Corollary 9. (Poincaré's Inequality) *For all special skew-symmetric tensor fields T in $\mathring{\mathbb{H}}(\text{Curl}; \Omega)$, i.e., for all functions $u \in \mathring{\mathbb{H}}(\text{grad}; \Omega)$ with $u \cong T$,*

$$\|u\|_{\mathbb{L}^2(\Omega)} \leq \widehat{c} \|\text{grad } u\|_{\mathbb{L}^2(\Omega)}.$$

Proof. We have $T \in \mathring{\mathbb{H}}(\text{Curl}; \Omega)$, if and only if $u \in \mathring{\mathbb{H}}(\text{grad}; \Omega)$. Moreover,

$$2 \|u\|_{\mathbb{L}^2(\Omega)}^2 = \|T\|_{\mathbb{L}^2(\Omega)}^2 \leq \widehat{c}^2 \|\text{Curl } T\|_{\mathbb{L}^2(\Omega)}^2 \leq 2\widehat{c}^2 \|\text{grad } u\|_{\mathbb{L}^2(\Omega)}^2$$

holds. □

We note that the latter Corollary also remains true for general skew-symmetric tensor fields $T \in \mathring{H}(\text{Curl}; \Omega)$ and vector fields $v \in \mathring{H}(\text{Grad}; \Omega)$ with

$$T = \begin{bmatrix} 0 & v_1 & v_2 \\ -v_1 & 0 & v_3 \\ -v_2 & -v_3 & 0 \end{bmatrix} \cong v.$$

§4. THE TWO-DIMENSIONAL CASE

Let Ω be a bounded domain in \mathbb{R}^2 with connected Lipschitz continuous boundary Γ , which is equivalent (in \mathbb{R}^2) to the topological property that Ω is simply connected. For tensor fields $T : \Omega \mapsto \mathbb{R}^{2 \times 2}$ we define analogously the Curl-operator row-wise by

$$\text{Curl } T = \text{Curl} \begin{bmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{bmatrix} = \begin{bmatrix} \text{curl } [T_{11} \ T_{12}]^t \\ \text{curl } [T_{21} \ T_{22}]^t \end{bmatrix} = \begin{bmatrix} \partial_1 T_{12} - \partial_2 T_{11} \\ \partial_1 T_{22} - \partial_2 T_{21} \end{bmatrix},$$

where now curl denotes the two dimensional scalar rotation and $\text{Curl } T : \Omega \mapsto \mathbb{R}^2$ is a vector. With the appropriate changes, Lemma 6 and Theorem 7 hold as well. In particular, there exists a positive constant c , such that

$$c \|T\|_{L^2(\Omega)} \leq \| \text{sym } T \|_{L^2(\Omega)} + \| \text{Curl } T \|_{L^2(\Omega)}$$

holds for all $T \in \mathring{H}(\text{Curl}; \Omega)$.

During the preparation of our paper we got aware that a two-dimensional related result may be inferred from Garroni et al. [4]. Instead of tangential boundary conditions $\nu \times P|_\Gamma = 0$ they impose the normalization condition

$$\int_\Omega \text{skew } T = 0. \tag{4.1}$$

Let us define the total variation measure of the distribution $\text{Curl } T$ for $T \in L^1(\Omega)$ by

$$|\text{Curl } T|_\Omega := \sup_{\substack{v \in \mathring{C}^1(\Omega) \\ \|v\|_{L^\infty(\Omega)} \leq 1}} \langle T, \text{CoGrad } v \rangle_{L^2(\Omega)}, \quad \text{CoGrad } v := \begin{bmatrix} \partial_2 v_1 & -\partial_1 v_1 \\ \partial_2 v_2 & -\partial_1 v_2 \end{bmatrix}.$$

We note

$$\langle T, \text{CoGrad } v \rangle_{L^2(\Omega)} = \int_\Omega T_{11} \partial_2 v_1 - T_{12} \partial_1 v_1 + T_{21} \partial_2 v_2 - T_{22} \partial_1 v_2.$$

Using partial integration, i.e., $\langle T, \text{CoGrad } v \rangle_{L^2(\Omega)} = \langle \text{Curl } T, v \rangle_{L^2(\Omega)}$ for $v \in \overset{\circ}{C}^1(\Omega)$, it is easy to see that $|\text{Curl } T|_{\Omega} = \|\text{Curl } T\|_{L^1(\Omega)}$ if $\text{Curl } T \in L^1(\Omega)$. In [4, Theorem 9] it is shown that for Ω having a Lipschitz boundary and a special ‘slicing’ property, there exists a constant $c > 0$, such that

$$c \|T\|_{L^2(\Omega)} \leq \|\text{sym } T\|_{L^2(\Omega)} + |\text{Curl } T|_{\Omega}$$

holds for all $T \in L^1(\Omega)$ with (4.1). Their proof uses essentially that in \mathbb{R}^2 the operators curl and div can be exchanged by a simple rotation, i.e., $\text{curl}[v_1, v_2]^t = \text{div}[-v_2, v_1]^t$. Thus, such a strong result may not be true in higher space dimensions $N \geq 3$ and it is open whether the normalization condition (4.1) can be exchanged with the more natural tangential boundary condition.

REFERENCES

1. P.G. Ciarlet, *On Korn's inequality*. — Chin. Ann. Math. **31B**, No. 5 (2010), 607–618.
2. J. K. Djoko, F. Ebobisse, A. T. McBride, B.D. Reddy, *A discontinuous Galerkin formulation for classical and gradient plasticity. Part 1: Formulation and analysis*. — Comp. Meth. Appl. Mech. Engrg. **196**, No. 37 (2007), 3881–3897.
3. F. Ebobisse, P. Neff, *Rate-independent infinitesimal gradient plasticity with isotropic hardening and plastic spin*. — Math. Mech. Solids, DOI: 10.1177/1081286509342269 **15** (2010), 691–703.
4. A. Garroni and G. Leoni and M. Ponsiglione, *Gradient theory for plasticity via homogenization of discrete dislocations*. — J. Europ. Math. Soc. **12**, No. 5 (2010), 1231–1266.
5. V. Girault and P.A. Raviart, *Finite Element Approximation of the Navier–Stokes Equations*. — Lect. Notes Math. **749**, Springer, Heidelberg (1979).
6. R. Leis, *Initial Boundary Value problems in Mathematical Physics*, Teubner, Stuttgart (1986).
7. P. Monk, *Finite Element Methods for Maxwell's Equations*, Oxford University Press, New York, (2003).
8. P. Neff, *On Korn's first inequality with nonconstant coefficients*. — Proc. Roy. Soc. Edinburgh Sect. A **132** (2002), 221–243.
9. P. Neff, K. Chelmiński, H.D. Alber, *Notes on strain gradient plasticity. Finite strain covariant modelling and global existence in the infinitesimal rate-independent case*. — Preprint 2502, <http://www3.mathematik.tu-darmstadt.de/fb/mathe/bibliothek/preprints.html>, Math. Mod. Meth. Appl. Sci. (M3AS) **19**, No. 2 (2009), 1–40.
10. P. Neff, D. Pauly, K.-J. Witsch, *A Canonical extension of Korn's first inequality to $H(\text{Curl})$ motivated by gradient plasticity with plastic spin*. — C. R. Acad. Sci. Paris, Ser. I **349** (2011), 1251–1254.

11. P. Neff, D. Pauly, K.-J. Witsch, *A Korn's inequality for incompatible tensor fields*. — Proc. Appl. Math. Mech. (PAMM) **11** (2011), 683–684.
12. P. Neff, D. Pauly, K.-J. Witsch, *Maxwell meets Korn: A new coercive inequality for tensor fields in $\mathbb{R}^{N \times N}$ with square-integrable exterior derivative*. — Math. Methods Appl. Sci. (accepted); <http://arxiv.org/abs/1105.5013>; Preprint SM-E-737, Universität Duisburg-Essen, Schriftenreihe der Fakultät für Mathematik, <http://www.uni-due.de/mathematik/preprints.shtml> (2011).
13. P. Neff, A. Sydow, C. Wieners, *Numerical approximation of incremental infinitesimal gradient plasticity*. — Preprint IWRM 08/01, <http://www.mathematik.uni-karlsruhe.de/iwrmm/seite/preprints/media>, Int. J. Num. Meth. Engrg. **77**, No. 3 (2009), 414–436.
14. D. Pauly, *Hodge–Helmholtz decompositions of weighted Sobolev spaces in irregular exterior domains with inhomogeneous and anisotropic media*. — Math. Methods Appl. Sci. **31** (2008), 1509–1543.
15. R. Picard, *On the boundary value problems of electro- and magnetostatics*. — Proc. Roy. Soc. Edinburgh Sect. A **92** (1982), 165–174.
16. R. Picard, *An elementary proof for a compact imbedding result in generalized electromagnetic theory*. — Math. Z. **187** (1984), 151–164.
17. R. Picard, *Some decomposition theorems and their applications to non-linear potential theory and Hodge theory*. — Math. Methods Appl. Sci. **12** (1990), 35–53.
18. R. Picard, N. Weck, K.-J. Witsch, *Time-harmonic Maxwell equations in the exterior of perfectly conducting, irregular obstacles*. — Analysis (Munich) **21** (2001), 231–263.
19. B. D. Reddy, F. Ebobisse, A. T. McBride, *Well-posedness of a model of strain gradient plasticity for plastically irrotational materials*. — Int. J. Plasticity **24** (2008), 55–73.
20. T. Valent, *Boundary Value Problems of Finite Elasticity*. Springer, Berlin (1988).
21. C. Weber, *A local compactness theorem for Maxwell's equations*. — J. Math. Anal. Appl. **2** (1980), 12–25.
22. C. Weber, *Regularity theorems for Maxwell's equations*. — Math. Methods Appl. Sci. **3** (1981), 523–536.
23. N. Weck, *Maxwell's boundary value problems on Riemannian manifolds with non-smooth boundaries*. — J. Math. Anal. Appl. **46** (1974), 410–437.
24. K.-J. Witsch, *A remark on a compactness result in electromagnetic theory*. — Math. Methods Appl. Sci. **16** (1993), 123–129.

Universität Duisburg-Essen,
Fakultät für Mathematik, Campus Essen
Universitätsstr. 2, 45117 Essen, Germany
E-mail: patrizio.neff@uni-due.de,
dirk.pauly@uni-due.de,
kj.witsch@uni-due.de

Поступило 14 декабря 2011 г.